Plukenetius

1691

41635/C/1

Vol. I (plates 1-120)

12908

LEONARDI PLUKENETIJ.

PHYTOGRAPHIA,

Sive

Stirpium Illustriorum, & minus cognitarum

ICONES,

TABULIS ÆNEIS,

Summâ diligentiâ elaboratæ;
Quarum unaquæq; Titulis descriptorijs,
ex Notis Suis proprijs, & Characteristicis
desumptis, insignita; ab alijs ejusdem
Sortis facile discriminatur.

PARS PRIOR.

MEMINISSE JUVABIT.



I Collins fec:

Ἔμπειρον. εἶναι συμβουλεύω ἰατρὸν εἰ δυνατὸν
πάντων φυτῶν. Gal. de Antidot. lib. 1. cap. 15.

LONDINI. MDCXCI.
SUMPTIBUS AUTORIS.

WELLCOME HISTORICAL MEDICAL LIBRARY

HONORATISSIMO

et

ÆTERNUM REVERENDO

PRÆSULI ac DOMINO

DOMINO

HENRICO COMPTON

LONDINENSI EPISCOPO &c. &c. &c.

Omnium Botanophilorum MECÆNATI summè colendo

LEONARDUS PLUKENETIUS

Has suas TABULAS PHYTOGRAPHICAS in
priori Parte comprehensas.
Quòd
HUJUS FAVORE ut plurimùm Illustrantur, submiſsâ cum
observantiâ, in devoti animi teſſeram grateq; mentis
symbolum, VOTIVAS reddit, jure meritoq;
Offert. Dicat. Consecrat.

54 e (2)

LECTORI BENEVOLO.

Natura parens sui studiosa conservatrix, artis & pulchritudinis suæ sibi conscia perditura fructum sui, si amplum hunc terrarum Orbem solitudini ostenderet, ut desiderium contemplandi fertiliſsimum stirpium apparatum; quem in ipsâ superficie subministrat nobiliſsima hæc quam pedibus calcamus Terra, in nobis excitaret & conservaret, Novitate animos nostros ad se allicit, dum indiès è varijs Asiæ Provincijs, & à Primoribus prorsùs incognitas in apricum producit plantas. Quamplurimi etiam Europæ loci à Majoribus nostris non ritè examinati Novas formas, Novasq species, quotidie nobis suppeditant. Africa quoq vel his nostris temporibus, monstra sua gignere non desijt: sed Novi aliquid ex Plantarum genere, semper ab extremis parturientis Africæ oris, ad nos adportatur. Nec hìc Natura subsistit, sed continuè Novas ex alieno Orbe, in lucem profert plantas. Novæq virtutes à filijs Artis, in salutem humani generis inveniuntur; Vigilant enim Sæculi nostri Botanici, qui (ut Cl. Raij verbis utar) nec Majorum inventis contenti, ad ulteriora tendunt, nec alios sibi inquisitionum terminos, & columnas figunt, quam quas Natura terris fixit, Universum Orbem curis suis complexi. Illi scilicet stimulo mihi fuêre, ut nulli sudori, operæ, ac sumptui parcerem, quin tam indigenas, quàm exoticas stirpes inquirerem, atq colligerem, quarum investigatio ab ipsâ Adolescentiâ nobis amæniſsimâ varietate se commendarit; fateor equidèm Botanica in omni ævo, melioris luti Quamplurimis excoli; et in Arte medendi vel antiquiſsima est ratio, per simplices Herbas, morbos abigendi; Hujus enim Cognitionem ad Principes, etiam & Magnates pertinere Salomonis (regum omnium quos Sol unquàm aspexit Sapientiſsimi) Alexandri magni, Mithridatis, Dioclesiani, & aliorum docuêre exempla; Imperatores & Reges non pauci, Heroesq Incluti, Indagationem, & peritiam hujus Naturalis Philosophiæ partis se digniſsimam judicarint, rati se Herbariæ rei colendo studium, optimè de humano genere mereri, nec minùs solidam, veramq laudem indè, quàm ex Triumphis, ad omnem Posteritatem, nomini suo libare poſse; Hanc tanti fecerunt Ethnici, ut ejus Inventionem, Dijs dignam æstimarint, & licet dubitare videantur, de primo eorum Inventore, consentiunt tamen tandem, Apollinem illâ prærogativâ maximè præeminere, his Poetæ verbis.

Inventum Medicina meum est, Opiferq per Orbem
Dicor, et Herbarum subjecta potentia Nobis.

et semper fuit adeò incredibilis apud Antiquos existimationis Herbarum Scientia, ut æqualis censebatur gloria, Herbam inveniſse & Vitam juvaſse; Quicquid enim hujus Sæculi sentiunt alij, nobis satis persuasum, Studium hoc Botanicum neceſsariam eſse Physicæ & Medicinæ partem, multis jucunditatis, & Delectationum condimentis gratiosam, & toti generi humano, utilitate quam maximè commendabilem: Visum igitur fuit, ut Nobiliſsimæ Botanices Scientiæ, ulterius inserviam, Plantarum aliquot Icones non alibi expreſsæ, periergâ manu, et emodulatâ ad Plantarum ipsarum dimensiones naturales, proportione quâdam, effigiare, & adumbrare: et earum plures, compendij gratiâ, in unam Tabellam conjeci; Opus non minùs varium, quàm ipsa est Natura, ubi Orientis & Occidentis Indiæ Montium, Sylvarum, Vallium, litorum & camporũ, Stirpes nativæ, & alumnæ à nullo Claſsico Autore anteà celebratæ, sed prorsùs Novæ, & hactenùs inauditæ, oculis vestris obversantur; singulis etiàm hujusmodi Novis plantis nomenclaturam proprijs & genuinis Symbolis insignitam accommodavimus, ut quæ rem ipsam aptiſsimè notat, & clarior foret omnis earum Cognitio simùl & Cognatio, quam insupèr diſsimilitudinis collationis & similitudinis ope, intèr Congeneres, & simillimas, iteratæ inspectioni, & ex Plantis ipsis expansioni, adaptavimus. Nam hæc omnia ex αὐτοψία didicerimus, non ut Nicander Colophonius, qui ab agro remotiſsimus, rustica aggredi, non veritus; neq ut Pamphilus, quem Galenus reprehendit, quòd cum non conspexerit; nè dùm per Somnium viderit, de his tamèn scripserit; Has nobis vidiſse contigit omnes, & etiamnum præ manibus sunt, quas in Herbario nostro vivo, imprœsentiarum conspicere licet, aliasq permultas, & indictas habeo, quas tamèn in præsenti non exhibeo, quià Chartæ penuria, & typorum immodica deterrent impensa; neq enim potest quis ullam certam cognitionem sibi arrogare Vegetabilium, quæ nunquàm aspexit. Benevolus interim Lector has Tabulas Phytographicas æquo suscipiat animo, donèc Herbarij vivi Catalogus, Plantarum circitèr Octo millium comprehendens consequatur, nec dubito quin candidi Rei Herbariæ cultores non solùm admiraturi sint, sed grati recolent, quòd tot incognitas (octingenti ferè numerantur) in his Tabulis nostris duas in partes distributis luci nunc damus, & si quid, vel etiàm Viris eruditioribus, unquàm Novitate suâ, se commendarit; Tabulæ nostræ Phytographicæ, Novi Orbis, Novis (i.e) noviſsimè detectis, & hucusq ineditis stirpibus, sunt refertiſsimæ, quibus amplum Exscriptoribus campum damus, in quo plantas legere poſsunt, easq generali Plantarum Indici, seu Pinaci inserere, quæ in nostro Orbe jamdudùm innotuêrunt; ex quibus omnibus non desperarem, nos Botanophilis perplacuiſse, & plurimas ab ijs gratias promeruiſse, tùm ob ea, quæ jàm præstitimus in hoc opere nunc edito, tùm ob ea, quæ præstituri sumus in Catalogo nostro, brevi edendo, ut hac ratione, votis meis ampliſsimè satisfactum iri arbitrarer. Utaris ergò Lector amice his nostris laboribus, quos amæno huic operi impendimus, ac stirpes hasce peregrinas rariſsimas, & hactenùs inauditas, quæ nunc exactis Imaginibus Orbi panduntur, adspicias, & in omnibus inexhaustam Sapientiæ profunditatem, & Majestatem divinam in ipsis Condendis comperies; His nobiscum ad Dei gloriam, tuamq, ac cæterorum utilitatem, & oblectamentum paratis, fruere pro otio, & incolumnis age; Vale.

"Plukenet is one of the [illegible] who [illegible] towards the end of the 17th Century. [illegible] who has given figures of [illegible] plants — has treated on the most scarce and rare plants in his Phytographia". [illegible] Elements of Botany.

"The indefatigable Plukenet procured and published an immense number of plants from all parts of the world, many of them very rare. His book is in every body's hands, and it would be superfluous here to say any thing of the utility [illegible] criticisms of his contemporary Plukenet have [illegible] but their value." Vid. Dr. Smith's Tracts p. 49

List of British Plants in this Volume, & the pages [illegible]

3.	Scirpus acicularis	40. [illegible]
	——— [illegible]	46. 4
	3 var	40. 5
	——— triqueter	40. 2
	[illegible]	40. 1
"	[illegible]	33.
"	Allium [illegible]	33. 0
"	Aira praecox	33. 4
"	Festuca ovina	34. 2
	——— [illegible]oides	33. 10
"	[illegible] junceum	35. 4
	[illegible]	37. 7
"	[illegible]	7. 5
4	[illegible]	75. 3
5	Convolvulus [illegible]	3 74. 3
"	[illegible] maritima	53. 8
"	[illegible] inundata	91. 3
"	[illegible]	62. 3
"	Ligusticum Scoticum	98. 2
"	E[illegible]	49. 4
9	Juncus [illegible]	46. 8
	——— [illegible]	32. 3
10	[illegible]	[illegible]
"	[illegible]	42. 8
13	Cistus [illegible]	[illegible]
"	Ranunculus [illegible]	[illegible] 1
	——— [illegible]	[illegible]
14	[illegible]	18. 3
"	[illegible]	41. 3
"	[illegible]	41. 1
"	[illegible]	[illegible]
"	Sibthorpia Europaea	[illegible]
15.	Arabis [illegible]	101. 3
17	Erodium maritimum	[illegible]
18	Fumaria lutea	[illegible]
"	Lathyrus [illegible]	[illegible]
"	Vicia sylvatica	[illegible]
"	Trifolium [illegible]	[illegible]
	——— [illegible]	[illegible]
	——— [illegible]	113.
"	Medicago maculata	113.
19	Senecio [illegible]	[illegible]
	var.	[illegible]
"	Hieracium [illegible]	[illegible]
"	Gnaphalium luteo-album	[illegible]
21	[illegible]	102.
"	Carex curta	[illegible]
	——— remota	34.
	——— [illegible]	[illegible]
	——— [illegible]	34. 7
	——— [illegible]	34.
	——— [illegible]	91. 8
	——— [illegible]	91. 7
23	Atriplex [illegible]	[illegible]
24	Polypodium [illegible]	[illegible]
"	Aspidium [illegible]	89.
"	Asplenium [illegible]	[illegible]
	——— Ruta muraria	3.
"	Pteris crispa	[illegible]
"	Cyathea [illegible]	89. 4
"	[illegible]	[illegible]
"	[illegible]	[illegible] 1
"	Sphagnum	101. 1
"	[illegible]	41. [illegible]
	[illegible]	[illegible] 1
"	Jungermannia	42. 2
	Bryum punctatum	45. 7



Tab. I.

1. *Abies Pinum referens folijs ex omni parte prorumpentib. Abies folijs prælong. Pinum simulans.* Ray. Hiſt. *an Picca minor* C B P. *Pinus maritima minor ejusd* Cat. H. Ae. Lugd.?

2. *Absinth. minus odoratū Gangeticum, floribus Chamæmeli solitarijs, e foliorum alis.*

3. *Absinth. tenuifol. elegans ex Hispania;* D.ris Turnefortij.

4. *Acacia ō Spinosa* Wagæ. Malabar. *Similis ē* Maderaspatan. *lit. a fol. ad justā magnitudinem denotat.*

5. *Acaciæ similis ō Spinosa ramulis alternis Maderaspatana. an Nili camarām. H. M. pro hujus Synonymo Canna Indicā arborea Zanon. ponitur à* Breyn. Prod. i. *cum Nellikam ejusd. hinc magis convenire videatur.*

6. *Acaciæ quodammodo accedens, S. Ceratiæ & Acaciæ media Iamaicensis spinosa bigeminatis folijs, flosculis ſtamin. atronitente fructu, ſiliquis parum intortis. lit. a. flosculos. b. Siliquā et fructum indigitant. Unguis Cati Arbor Americana spinosa ſiliquosa* Par. Batav. Prodr.




Tab. II.

f. 1. Acacia Maderaspat. ſpinosa Intsiæ accedens, cortice cinereo, ramulis communi pediculo binis. a. folium ad natural. dimensionē exprimit.

2. Acacia gloriosa Lentisci fol: ſpinos. flore spicato luteo, ſiliquâ magnâ muricatâ. Arbor exot. ſpinos. fol. Lentisci C.B.P. Inimboy Bras. Marcgr. Bonduch Zanon. Caretti H.M. lit. a. folium ad verā magnitud. cum aculeo sursûm ſpectante. lit. b. lobum Spinis crebriorib; armatū indicat.

3. Acacia ō spinosa minor ē Quaracoa ſ. Curaſsavica fl. alb. globoso. fortē Tehoaxin Mexicens. Hernand.

4. Acer virginianum folio majore ſubtùs argenteo, ſuprà viridi ſplendente. Acer floresc. Virginiana Vulgō. Arborē hanc magnitudine & proceritate eximiam in Hort. D. Ioh. Tradescanti Lambethæ ad Austrū jamolī enutritam ſæpè conspeximꝰ sed eam ibi unquàm floruiſse a nemine memoratum audîjmus.



Tab. III.

f. 1. Acer Benghalense Laurin. folijs fructu tergemino. totidèm membranulis extantib; alato.


2. Adianth. alb. floridũ. s. Filicula petræa crispa Ray Hiſt. *Adianth. album crispũ Alpinũ Schwenckfeld.* I.B.

3. Adianth. folijs minutim in oblongum ſciſsis. pediculo viridi. C.B.P. *Ad. alb. tenuif. Rutæ murariæ accedens* I.B.

4. Adianth. album florid. Cicutariæ folio Virginianũ nostrum. D. Ray Hiſt. app.

5. Adiant. radicos. humisparsũ. s. Filicula pellucid. nostras Coriandri foliolis mollicellis globuliferum Adiant. petræum perpusillum Anglicũ folijs bifid. vel trifid. Newtoni Ray. Hiſt. *forte Filicula saxatilis fol. Coriandri.* Mentz. pugil.

6. Adianth. radicos. erectius foliolis imis biſectis, cæteris verò in-tegris tenuiſsimè crenatis. forte Filicula *saxatil. Coriandri folijs rotund. leviter serratis Mentzel. pugil. Rarior. Pl.*



Tab. IV.
f.1. Adianth. nigrum Chinens. tenuitèr diviſum, pinnulis minimis, obtuſis, plerumq; bifidis. Ray. Hiſt. app. an Filix ſaxatilis criſpa Park? lit. a. ramulum ad juſtam magnitudinem denotat.

2. Adianth. nigrū ſ. Dryopteris Chinenſis lanuginoſa Ray. Hiſt. app. an Lonchitis Maranthæ, Ceterach ramoſum Quorund. Lob. Ic. ſ. Polypodium Iluens. Hiſt. Lugd. quæ ſecundùm utroſq; Bauhinos ſunt unū & idem:

3. Adianth. Chinenſe perelegans ramoſ. folio flabelliformi cum rubedine perfuſo. Adianthum folijs Coriandri ramoſum Ray. Hiſt. app. has tres Adianthorum ſpecies habuit Cl. Rajus ab. egregio Botanico & Pharmacopæo Londinenſi D.no Iacobo Petiver qui poſteà ad Icones exſculpendas, nobis libentèr conceſſit.

4. Æſchynomene mitis ſ. Mimoſa De Tunquin pinnulis amplioribus Chamæfilici marinæ æmulis. ſub Ovali titulo recepimus, foliorum formâ Tſjampangam H. M. haud multùm abludit.



Tab. V.
f. 1. Æſch. mitis Americ. foliatis Acaciæ brevioribus; Br. Cent. 1 Mimosa Americ. supina veluti spinulis exasperata, pinnis Acaciæ latiorib; & brevioribus ejusd. Prodr. 2.


2. Æschyn. mitis ſ. Mimosa angustiori siliquâ folijs perexignis Maderas-patana Æsch. folijs minimis nostra. Ray. Hist. app.

3. Alcea Indica frutesc. folijs ad marginem exasperatis Brijoniæ albæ divisuris. è Sinu Bengalens. allata est nobis.

4. Alcea Spinōs. Maderaspatana fol. profundè laciniatis, Cannabis in modum penta-phyllæa. ſaporis acidi an Acetosa Indica Bonti. Huj; meminit Cl. vir & ſedulus exoticar. Pl. explorator D^r. Herman in Cat. H. Ac. Lugd. pag. 27.



Tab.VI.
f.1. *Alcea Cretica flore immaculati candoris peramplo. in Hort. Medic. D. Morgan Londini ſubur bano, jam olim exculta fuit.*
2. *Alcea acetoſa Indica mitis folijs ſuperioribus Goſſipij in modum tripartito diviſis. Althæa Indica Goſſipij folio Acetoſæ ſapore* CHAL.
3. *Alcea Indic. Ulmi folio prægrandis subtus incana flore amplo purpureo, Althæa magna 5q. capsularis folijs integris subtus albicantibus, floribus magn.* Banist. Cat. Virgin. apud Ray Histor. *inferiora enim hujus folia ſunt prorsus integra, ſuperiora tamen in lacinias ternas leviter inciſa, & aversâ parte albicantia.*
4. *Alcea ſ. Bamia American. parvo flore, petalis rotundioribus.*
5. *Anchusa lutea minor* C. B. P. I. B. Park. *item Symphytum angusti fol. Apulum* ejusd. Park. *Anchus. echioid. lutea Cerinthes flore, montana.* Col. ecphr.



Tab. VII.

f. 1. Alcea Ind. fol. hastato, fl. parvo, purpur. longo petiola insidente, pericarp. compacto, duro in 5 qq. cellulas, totidèm semina includentes dispertito.

2. Althæa frutesc. Ind. Orient. Populi albæ minoris fol. tenniter dentato molli. an Althæa Americ. frutesc. Populi alb. folijs Par. Bat. Pr.

3. Alsine Caryophylloid. tenui fol. fl. albo punctato. an Als. Coryophylloid. tenuifol. montana Lini flore Ray Hist. fortè etiam Polygonum Caryophylli Holostei flore Cat. H.R. Bles. Betonica. s. Tunicæ minimæ similis Planta I. B.?

4. Alsin. palustr. Ericæ folia Polygonoides, articulis crebrioribus, flore albo pulchello. Saxifraga palustr. Anglica Goodyeri Park. item Alsin. nodosa Germanic. ejusd. Park. Alsine nodosa Germanic. aliàs Arenaria. C.B.P. Arenaria I. B.

5. Alsine formis paludosa tricarpos flosculis albis inapertis. Alsine palustris Portulacæ aquatic. similis. Ray. Hist. Alsin. palustr. minima flosculis albis, fructu Coriandri exiguo Mentz. pugil. Portulaca exigna. s. arvensis Camerarij I. B.

6. Alsin. spuria pusilla repens folijs Saxifragæ aureæ Ray. Cat. Pl. Angl. Hujus notitiam Doctiſs. nostro Raio debemus, proindè miratus sum quo casu in Caleberrim. Autoris Histor. Universal. hæc plantula omiſsa fuit.

TAB. VIII.



Tab. VIII.

f. 1. *Althæa frutesc. folio acuto virente molli, flore specioso, an Althæa frutesc. folio acuto flore magno* Schol. Botan. *Althæa profundè serrato. s. dentato folio* I.B. *Althæa flore majore* C.B.P. *et fortaſsè eadem cum Althæa arborea Olbiæ Provincialis* Advers. Lob. & Icon. *Malva sylvestr. perennis folio profundè laciniato & dentato* Moris. Hist.

2. *Althæa frutesc. Bryoniæ folio incana flore minore. Althææ species quibusdam arborea altera, vel Bryoniæ folio* I. B. *Alth. sempvirens Bryoniæ fol.* Park. *Alth. frutesc. Bryoniæ folio; itèm Alth. frutesc. folio acuto parvo flore* C.B.P. *Malva frutesc. folio acuto auriculato Hederæ instar* Moris. Hist. *Althæa Olbiæ vulgò putata.*

3. *Althæa fruticans Hispanic. Aceris Monspeſsulan. incanis folijs grandiflora saponis odore. Alth. frutesc. folio rotundiore incano* C.B.P. *Alth. arboresc.* I.B. *Alcea frutesc. folio rotundiore incano* Moris. Hist. *item Malva Hispan. folijs mollib; undulatis in margine superiùs micis sulphureis ad solem lucentibus donatis ejusd* Moris. Hist. *quoād descriptionem; at in Tabul. huic Titulo alterius Icon substituitur, Althæa Provincialis* Penæ Advers. *ex sententia* D. Rāy Hist. 603.



Tab. IX.

f. 1. Alsines facie myosôtis Bisnagarica frutescens flosculis pallidè flavescentib; cujus folia sunt ad caulem alternò positu, (non per intervalla geminata) incana, hirsuta; flosculi quidèm ut Anagallidis, non tamèn cœrulei, alio quin haud admodum diversi nimis incautè, est enim reverâ Convolvuli minimi species Bisnagarica.

2. Althæa Indica latiori folio cordiformi, ad summum sinuato. Malva Indica frutescens folio cordato P. B. P.

3. Althæa rostrata Cormandeliens. Pimpinellæ major. fol. subrotundo.

4. Althæa fruticosa minor Gangetica folijs exiguis margine spinosis.

5. Althæa Ind. floscul. parvis folliculatim (fortè fasciculatim) ramulis adfixis. Breyn. Prodr. i. Malva American Carpini folio, florib; aureis fasciculatis parvis P. B. P. è Maderaspatan nobis allata est.

6. Althœa Virginian. bidens Pimpinellæ maj. acutiore folio, flosculis minimis luteis. fortè Alcea Mori folio Virginian. min. Sch. Bot.



I. Collins. de: & fe.

Tab. X
f. 1. *Amaranth; albus Gangeticus Salviæ angustis folijs paniculâ conglomeratâ. an Amarant. Alopecuroid. Indicus frutesc. lanuginos; folijs Salviæ angustiorib; & longiorib;* Br. Prod. 2? ***Salvia Amaranthoides Mich. Nutij Fascic. Herbar.***

2. *Amarant. Ind. min. lanuginos. Sampsuchi folijs, spicâ albâ ad foliorum exortum geniellâ. an Amarant. Alopec. Ind. frutesc. lanuginos; folijs Salviæ latiorib; & breviorib;* Br. Prod. 2?

3. *Amarant. verticillat; min. Bengalens. Serpijll. folijs incanis.*

4. *Amarant. spicat; Dictamni Creticæ folio Maderaspatenſis.*

5. *Ambrosia maxima inodora Marrubij aquatic. folijs tenuitèr laciniatis Virginiana.*

6. *Ambrosia (forsàn) è Chinâ Anguriæ folijs accedens. à D.no Petiver recepi.*



1. *Ammi majus exoticum folijs per ambitum eleganter inciſis. an Ammi alterum semine Apij* C.B.P. *Ammi odore Origani* I B. *nostra folia à Dauco Cretico* Zanon. *haud multùm abludunt.*

2. *Amomum Virgin. Evonymo Virgin. simile, florib; & fructu sparsim erumpentib; Amomum Virginianum Cornifæminæ facie* Ray Hist.

3. *Amygdalus Indica nana.* Cat. H.R. Paris. *Amygdalus pumila* Cat. H.R. Bles.

4. *Amygdalo-persica nana flore carneo pleno Africana. Persica Malus Africana nana flore incarnato pleno* Cat. H.Ac. Lugd. *hujus ramulum. a. floribus onuſtum denotat.*




Tab. XII
f. 1. *Angolam fortè* H. M. *Arbor Indica baccifera fructu umbilicato, rotundo, Cerasi magnitudine, dicocco.* Ray Hiſt. *ē sinu Gangetico nobis advenit.*

2. *Anisum Africanum fruticeſcens folio, & caule, rore cæruleo tinctis. Ferula fruticosa sempervirens folijs Anisi Galbainfera, ex quâ Galbanum Officinarum* P.B.P.

3. *Anthyllis s. Alsine maritima lignosior Anthylloides, folijs tomentosis Bisnagarica.*

4. *Antirrhinum purpureum multicaule, folijs tenuiſsimis, sursūm inflexis, florib; ſparſis, calyce spinulis asperato.*



Tab.XIII.

f.1. *Apocynum Canadense maximum flore minimo herbaceo* Cat.H.R.B.les. *itèm Apocyn. Virgin. flore herbacco, ſiliquâ longiſſimâ, & tenuiſsimâ* Moriſon. Prælud. Botan.

2. *Arbor Ariſtolochiæ folijs Maderaspatana fructu parvo, Medicæ adinstàr involuto. an Gummi Elemi fundens Arbor Orientalis?*

3. *Arbor baccifera Orientalis, folio rhomboide, fructu calyculato. Caſsia Cinamomea s. Canella de mato.* Linschot. part. 4. Ind. Orient.

4. *Arbor è Cormandèl vasculifera folijs amplis Mijrsinites rigidiorib; nostra* Ray. Hist.



Tab. XIV.

f. 1. Arbuscula Bengalensis, Aceris minore folio incano alternatim posito.

2. Arbuscula è Maderaspatan folijs Aceris alternâ vice lucidè virentibus.

3. Arbuscula Zeylanica Aceris folio baccifera, folijs è directo, magis angulosis.

4. Arbuscula Bisnagarica Aceris folio parvo aculeata, folijs è regione binis

5. Arbor aromatica è Cormandel Laurinis folijs binis ternisve. nostra D. Ray. Hiſt.

6. Arbor Prunifera. sphærulas saponarias ferens tetraphylla, ex Ind. Orient. an Saponaria Arbor Zeylanic— trifolia semine Lupini: Conghas Zeylanensib; P. B P.

7. Aster Americanus frutescens Satureiæ folijs scabris floribus amplis, saturatè Violaceis.



Tab.XV.
f. 1. *Artemisia Chinensis cujus mollugo Moxa dicitur, fortè Ytzecuinpatli Mexic.* Hernand. *an Artemisia Virginiana* Park?
2. *Aristolochia Violæ fruticos. folijs virginiana, cujus radix Serpentaria dicitur.*
3. *Asarum è Terrâ Mariana Violæ luteæ vel Asarinæ folijs, Serpentaria nigra Nostratib; dicta.*
4. *Asparagus aculeatus minor sarmentosus è Maderaspatan.*
5. *Asparagus (fortè) aculeatus major spinas plurimas ab eodem exortu pro folijs ferens. an Asparag; aculeatus Siculus Raij Hist.? an Spina umbellâ folijs viduâ C. B P.? Drypis Theophrasti Anguillaræ I B?*
6. *Aster exoticus* Hyoseridis *folio minore noster* D. Ray *Hist.* app.
7 *Aster African; Stæchadis inodoræ folijs incanis flore albicante. Aster Afric. Stæchad. folio noster* Ray H. app
8 *Aster Bengalensis omnium minimus, summo caule biflorus.*



Tab. XVI.

1. *Aster luteus latifol. glaber folijs rigidis, & minutiſsimè crenatis. an Aster luteu: folio glabro & crenato CBP.? ubi Synonyma huic Plantæ à Casp: Bauhin. malè sunt appropriata.*

2. *Aster peregrinus Cisti folio non crenato, flore magno luteo. an Aster odoratus flore luteo stellato exoticus* Cat. Pat. *à Turre: fortè. Aster Conyzoides luteus perennis* Icon Roberti.

3. *Aster luteus fœtidus Crithmi Chrysanthemi folio, florib; umbellatis. Iacobæa Crithm. litorei folijs* Catal. H.R.Bl. *Iacobæa fœtida Crithmi Chrysanth. facie Bruniyeri.* Cat. H. Ionq. *Iacob. Rosmar. fol. Quorund.*

4. *Atriplex maritima Mauritanica frutescens, Polygoni folijs argenteis racemosa. Atrip. marit. Mauritan. fol. Polygon. argent. semine amplo fusco vasculis rotundiorib; nostra D.* Ray Hist. append.



Tab. xvii. f. 1. Bellis montana incana & crenata, folijs & crenis obtuſis. an Bellis montana folio obtuſo crenato. C B Pin & in Prodr.? Bellis sylvestris folio craſſo & villoso. Sch. Bot.? ubi ponitur pro nondùm descripto.

2. Bellis Americana procerior serotina ramosa flore ampliſſimo Cat. H. Ac. Lugd. an Bellis radice repente, folijs latiorib; serratis. Cat. H. R. Bles. fortè Tlally-yetl. Nov. Hispan. Hernand.? Bellis Indica maxima C. H. Ac. Pat. Turre.

3. Bellis montana folijs Coronopi lacinijs s. denticulis rarioribus & magis extantib; an Bellis Alpina folijs profundè diſſectis Sch. Bot.? Bellis leucanthemia (s. Anthemis) fruteſcens folijs profundè laciniatis Breyn. Prodr. 2. Bellis montana folijs Chrysanthemi D.ris Magnol ejusd Breyn. ibid. Bellis chrysanthemi segetū folio D.ris Hans Sloan. Ray Hist.

4. Bellis pumila crenata Agerati æmula, crenis bicornibus asperiusculis. Bellis maritima folijs Agerati C B P & in Pr.

5. Bellis montana Tanaceti folijs caule singulari annua. Chamæmelo affine Buphthalmum Italicum segetum altiſſimum I B.

6. Bellis Alpina Parthenij folijs elatior.



Tab. XVIII.

f. 1 Biti Malabaricæ accedens Arbuscula Maderaspatana flore peculiari.

2 Blitum Africanū Calthæ palustris folio caule nudo cubitali spicam peda=
=lem & ampliùs sustinente. Petasites Africanus Calthæ palustris folio Cat.
H. Ac. Lugd. Petasitidis folio Bliti fructu D. Ray Hist append.

3 Bugula cærulea Alpina Park. Consolida media cærulea Alpina C B P. &
I B. item Consolida media Genevensis ejusd I B. Histor. Universal.



Tab XIX

f. 1. Carduus Andelusiacus spinoſiſsim; echino grandi
aculeis rigidis armato noster D. Ray Hist.
append. é Regno Andelusiæ, (ubi & ipse
collegit) hanc elegantiſsimam Cardui
speciem mittebat ad nos Ornatissimus Vir
& nobis amiciſsim; D. Ioh Fishwick.



Tab. XX.
f. 1 *Christophoriana arbor aculeata Virginiensis. Angelica arborescens spinosa s. Arbor Indica Fraxini folijs cortice spinoso* Ray Hist. *Arbor spinosa Fraxini folijs Virginiana.* D. Edward. Morgan. *Angelica arbor vulgō. lit. a. folium hujus, naturali magnitudine duplô minus. b. seminum Umbellas indicant.*



Tab.XXI.

f.1. Chryſanthemum Americanū Doronici fol. fl. Perſici coloris, umbone magno prominente, ex atro-purpureo, viridi, & aureo, fulgente, an Bellis Canadenſis purpurea Ciaſsi Fascic. Herbar. Mich. Nuty p. 22.

2. Chryſanthem. Canadens. latifol. humilius, Moris. Præluд. Bot. Chryſanth. Canadens. humilius. Boccon de Pl. Sicul. Aster virgineus luteus alter minor. Park.

3. Chryſanth. African. Aſteris facie, imo flore non folioſo, capitulis duris.

4. Chryſanth. Bengalenſe auguſti-foliū puſillum. ſummo caule ramoſum.

5. Chryſanth. ſegetum noſtras folio glauco multiſciſſo maj; fl. minore.

6. Chryſanth. tennifol. flore luteo subtūs purpūraſcente. Buphthalmū. tennifolium folio Millefolij ferè I.B. Buphth. caule & folio purpuraſcente C.B.P. Buphth. flore purpuraſc. Park.

7. Chryſanth. Smyrnæum Cotulæ folio, flore amplo nudo ſeminib; hiſpidis, ad capitula in orbium ſpiras circumvolutis. an Chryſanth. Valentinum, Clus (ie) Buphthalmum lanuginoſum folijs Millefolij. C.B.P. Elegans hæc planta cujus folia aliquandō lanugine obducta invenimus ex ſemine in Cabis s. Corbulis quibus Ficus & Uvæ paſſæ è Smijrnâ ad noſ transportantur, diſperſo & indē collecto, enata, Angliæ primò & noſtræ demum notitiæ acceſſit.



Tab XXII

f. 1 Chryſanth. Scrophulariæ folijs Americanum.

2. Chryſanth. Cannabinum Virginian. hirſutum, diſco nigro, petalis aureis radiato. Chryſanth. s. Flos ſolis laciniatus annu; umbone nigricante D. Bobart an Eupatorium Cannabinum Americanum flore aureo Cat. Hort. Reg. Bles. lit. a a. ſummum ramulum ſuis floribus ornatum denotant.

3. Chryſanthemum Peruvianum flore albo Cat. Hort. Lips. olim. edit.

4. Chryſanth. Chinenſe folijs plurifariam diviſis. Halicacabi peregrini æmulis; Chryſ. fol. multifidis Cicutæ nonnihil ſimilibus Virginianum Cat. H. Ac. Lugd. Chryſ. Cannabinum American. frutesc. Virginianum Sambuci laciniatæ folijs & facie Breyn. Prodr. 2. è Chinâ ad D. Petiver allata est hæc elegans planta, & ab illo nobis amiciſſimè communicata.



Tab. XXIII.

f. 1. *Campanula Cymbalariæ folio Park. Camp. Cymbalar. fol. vel folio hederaceo C.B.P. Campanul. folio hederac; species* Cantabrica Anguillaræ. I B.

2. *Cicer Astragaloides (forte) virginianum, hirsutie pubeſcens florib. amplis, subrubentibus. an Cicer montan. lanuginos. erectum* C.B.P. D Banister Cat. Stirp. Virgin.

3. *Cistus American. Chamænerij folijs lucentib;* ex Hort. D. Morgan. *an Tzatzoqua Mexicens.* Hernand.

4. *Cistus Chamærhododendros folijs confertis, ferreà rubigine nitentib; margine pilosis.*


5. *Cistus Helianthemos argenteus folio latiore, flore saturatè luteo. Helianthem. Alpin. folio Piloſellæ minoris Fuchsij IB. ex sententiâ* D. Raj

et forte Cistus helianth. folio Polij montan. ejusd. *item Cistus pusillus folio candido, flore luteo,* ejusd. *IB. ex mente* ipsius Raÿ. Hist.

6. *Cistus humilis Alpinus durior, Polij nostratis folio candicante, Chamæcistus montan; Polij folio noster* D. Raÿ. Synops. Stirp.

Brit. *an Chamæcist, folijs Myrti Tarentin. canis & cinereis IB.*

7. *Cisti Helianthemi folijs, planta Maderaspatana, floribus ex alis. an Vitis Idæa Cisti helianthemi folijs Maderaspalana.*



Tab.XXIV.

f 1. *Clinopodium birginianum angustifoliū. fl. amplis luteis, punctis purpureis: Clinopod. Virgin. anguſtif. flore luteo* Cat. Hort. Ac. Lugd. *Origanum floribus amplis luteis purpuro-maculatis, cujus caulis sub quovis verticillo decem vel duodecem. foliolis rubentibus est circumcinctus.* D. Baniſt. Cat. Stirp. Virgin.

2. *Conferva reticulata crispa.* D. Doodij Ray Hiſt. app.

3. *Convolvulus anguſtiſsimo folio nostras cum auriculis. an Convolu. arvensis minim;* D. Dent Ray. Synops. ſtirp. Brit.

4. *Convolvulus maritim. s. Soldanella* è Maderaspatan; *folio in summitate sinuato. an Convolv. marit. Zeylanic; folio crasso cordiformi* Cat. H. Ac. Lugd. *Bintamburù Zeylan.*

5. *Convol. maritim; majore folio Chinensis. an Sarsa litoralis. s. Convol. maxim; Braſilieſ. Pison.*

6. *Convolvuli instar Volubilis* è Coromandel *umbilicatis folijs craſsis acuminatis, villosa. an Pada-Valli, s. Pada Kelengi sive Muschega-Pada H M. P. 7?. fortè ad Brijoniæ genus, potiori jure releganda.*



Tab. XXV.
f. 1. Convolvulus American; sub Ialapiæ nomine receptus. an Convolvul; Colubrinus Piſon? Caapeba s. Convolvul; Braſilian; flore octapetalo monococcos Raÿ. Histor?
2. Convolvul; s. Volubilis Americ. Bryoniæ nigræ nervoſis fol. vasculis seminiferis, triquetris. Helxine fruticeſcus. Bryoniæ nigræ folijs, capsulis triquetris amplis, pergamenis. D. Banist. Cat. stirp. Virgin. an Capreolata Bryoniæ nigræ folio Braſiliens. tricocca Marcgrav.
3. Convolvulo similis Bryoniæ albæ folijs villosis, ex India Orientali. an Nehoemeka H. M. P8. Tab 19? Hæc autem pro synonymo Bryoniæ Zeylan. folijs profundè laciniatis Cat. H. Ac. Lugd. à Cl. Comelino ponitur. Annot. in loc.
4. Convolvulo similis villosa, folijs craſsis, subrotundis, è Maderaspatan.



Tab. XXVI

f. 1. *Corallina latifolia & Opuntia marina* Cortus. *Fucus folio rotundo* C.B.P. *Sertolaria* Imperati, *Scutellaria s. Opuntia marina* I.B.

2. *Corallina fistulosa fragilis, internodijs prælongis, lævibus, albis, Farciminum modo catenatis, Corallina fistulosa fragilis* I.B.

3. *Cornus virginiana flosculis plurimis albidis ex involucro tetrapetalo rubro erumpentibus*, D. Banist. Cat. Stirp. Virg. apud D. Ray Hist.

4. *Cucumis Bryonoides Bisnagarica fructu parvo, florum calyce muricato. an Mucca Peri.* H.M.P. 8.

5. *Cuscuta ramis Arborum innascens* Caroliniana *filamentis lanugine tectis*, Musæi Courteniani.

6. *Cuscuta lendiginosa tenuiſsimis cirrhis* è Terrâ Marianâ. *Cuscuta Americana super arbores se diffundens*, Ray Hist. app.

7. *Cytisus Americanus flore luteo spicato; folijs suprà virentibus, glabris, subtùs lanugine argentatis.*



Tab. XXVII.
f. 1 Elichryso affinis Peruviana frutesc. Cat. H. Ac. Lugd. app. Agerato affinis Peruviana frutescens Parad. Bat. Prod. Quinquina. s. Arbor amara & Febrifuga quibusd. falsō & fastuose nuncupata.
2 Elichryso affinis virginiana frutescens folijs Chenopodij glaucis. Senecio frutescens virginiana Atriplicis folio Raij Hist. & app.
3. Ephemerū Bengalense serpens folio subrotundō brevi phalangoides. an Ephemerum Africanū annuum flore dipetalo Cat. H. Ac. Lugd. Dipetalos virginiana folijs Gentianæ, aut Plantaginis Marcgr. Raij Hist? fortē etiam Gentianella Indica. Bontij. p. 154.
4. Ephemerū Phalangoides Maderaspatens. minimum folijs perangustis perfoliatum, fortē Acaxaxàn. s. Potamogeiton Mexiocan. 2. Hernand. Planta palustris & lenta, quæ statim comprimentib; cedat digitis, huc etiam pertinet Tali-pullu. H. M. P. 9. T. 63. p. 123.
5. Euphrasia lutea latifolia palustris. Raij Hist. an Cristæ Galli affinis planta Romana, s. Crista Galli major Italica I. B. quærit & ipse Eruditiſſ. Rajus ubi supra.



Tab. XXVIII

f. 1. *Euphrasia. s. Cratæogonum folijs brevibus obtusis Vestmorlandicum* Ray Hist.

2. *Euphrasia pratensis minor purpurea* Park. *Euphr: purp. min.* C.B. Prodr. *Euphr. 3.ª non scripta* Fab. Col. ecphr.

3. *Ericæ facie Polygonoides summitas cum floribus.*

4. *Ericæ facie Polygonoides Africana pentapetalos, viticulis in creberrima capilla-menta squammulis conflata diffusis, Limonium feru-laceo folio,* Grisley Virid. Lusit. *ut humaniſsime nos monuit D. Gulielm;* Sherard. *Collegij S.ti Johannis in Academ. Oxo-niens. socius, & Phytognostes peritiſsimus, cujus munere summitatem accepi, ramulum autem à D.* Iohanne Fish-wick. *ad maritimas Africæ oras à seipso collectum & ad nos transmissum, obtinui.*

5. *Evonymus virginian; rotundi fol. capsulis coccineis eleganter bullatis* Banist. Cat. Stirp. Virg. *lit. a. a. semina ad natural. magnitud. demonstrant.*

6. *Evonym. Jujubinis folijs Caroliniensis, fructu parvo ferè umbellato. b. b. semina ad æqualem dimensionem significant. à foliorum simili-tudine cum Rhamno Catharctico (ni fallor) hæc planta, Spina Cervina Barbadensibus nostratibus nominatur.*



Tab. xxix.

f. 1. *Ficoides s. Melocactos mamillaris glabra, sulcis carens, fructum suum undiq; fundens. an Echino-melocactos minior lactescens absq; tomento, cylindris laxioribus Parad. Bat. Prodr.*

2. *Ficoid. s. Melocactos minim. lanuginos; spinis mitioribus fructu sparsim egrediente. an Echino-melocactos tuberosus, s. globularis min. Cat. H. R. P.*


3. *Ficoid. Americanũ. s. Cereus erectus cristatus folijs triangularib; profunde canaliculatis. Cereus spinis crebrioribus horridus Ray. Hist. Jamacarũ prima Pison. Hist. Lusitan. Cardon. an Planta pinnata arundinacea Park. Theatr.*

4. *Equisetum s. Hippuris lacustris, folijs mansu arenosis Gesnero Ray. Hist. Hippuris coralloides Ger. emac.*

5. *Frutex Cormandeliensis baccifer Mali Granatæ folijs amplioribus, noster Ray. Hist. 1801.*

6. *Frutex Gangetic; Alaterni folio spinosus. Frutex aculeatus Orientalis Ilicis folio noster Ray. Hist.*

7. *Frutex Bisnagaricus spinosus, Coryli folijs densis, spinis brevioribus crassis.*



1. Filix. ampliſsima lobis foliorum lacinialis Cambrica Polypodium Cambrobritannicum pinnulis, ad margines lacinialis Raij. Hist.
2. Filicis ē Chinâ genus molle, auriculâ ad pinnulæ basim supernē productâ, summo folio longiūs mucronato.
3. Filix. s Polypodium Indianum folijs profundē sinuosis Marrubij aquatic. æmulis. an Filix Indica Osmundæ facie Bod. ā Stapel. in Theophrastū Sim. Paul. Quadripart. Botan. 302.
4. Filicifolia, Lonchitis aspera maxima Indorum.



Tab. XXXI

F. 1. *Frutex baccifer. e. Cormandel Roriſmar. obtuſis & veluti punctat. fol. plurimis, ex uno tuberculo provenientib. vid.* Raij Hist. p. 1801. *an Vitis Idææ spec. Indicæ Orientalis.*

2. *Frutex baccifer folijs Lini, vel Caſsiæ Poeticæ Maderaspatensis. noster.* Raij Hist.

3. *Genista tinctoria Maderasp. Roriſmar. folijs ad caulem radiatis. an Legumen cujus siliquula à D. Terentio ex Indiâ missa, & titulo herbæ Anil inscripta fuit. C.B.P. Glasto affinis ejusd. C.B.P. Anil s. Nil, Indorum color. I.B. Anil herba folijs Mungiriqua similis Garziæ: Anil s. Indigo, Gali, s. Nil, Herba Roriſmarini facie, Linschot. part. 4. Ind. Orient. Genistæ Iluensi Lugd. (h. e.) Genist. radiatæ, vel stellari I.B. (i. e.) sparto triphyllo C.B.P. baldè similis, si non eadem. Huic etiam proxima est Cytisus, acuminatis folijs, C.B.P. quoàd D. Turnefortiū monente D. Sherard (i.e. nostrâ sententia) Cytisus Indicus flore luteo hirsutus Cat. Patav. à Turre. Cytisus folijs argenteis flore luteo vulgò. Nostræ caulis angularis est, proindè eam ad Genistarum genera magis propriè disponendā arbitramur; Quam ut audio præparant prô cœrulei coloris tincturâ in Regno Bisnagar, alijsq. in locis Indicæ Orientalis.*

4. *Geraniū pusillū supin. marit. Alth. aut Betonic folio nostras* Raij Hist. *Geran. minim. procumbens folijs Betonic.* Moris. Hist. 2.

5. *Gnaphal. Lusitan. comis argenteis. an Chrysocome Syriaca candidiss. florib. Breyn. Pr. 1. fortè etiam Gnaph. maj. angust. oblongo folio alterum C.B.P. Filago Germanic. altera secundùm folia florida, Lob. Icon.*

6. *Gnaph. maj. lato oblongo fol. C.B.P. Gn. 2m. Plateau. Clus. Hist. Gnaph. ad stæchadem. citrin: accedens I.B. Elichryson silvestre latifol. capitulis conglobatis* C.B. Phyt.

7. *Gnaph. impium s. Filago dicta Cap. Bon. spei. Gnaphalio Plateau 3o. simile genus, sed diversum.*



Tab. xxxii.

f. 1. Gramen Arundinaceum Halepense, Tragopogonis fol. paniculâ miliaceâ, huj; exemplar eximij Botanici D. Iac. Bobart. Hort. Academicæ Oxon. Præfect. communicatione in Herbario nostro vivo servatur.

2. Gram. paniculat. Gallo-provinciale Aquilegiæ semine Schol. Botan. a D. Gulielmo Sherard accipiebam.

3. Gr. junceum aquat. paniculis cum fol. capillarib; simul ortis, proliferum. an Gr. junc. capsulis triangul. caudicul. tenuib; folijs ad nodos, & panicularum divaricationes prælongis, Ray Hist.

4. Gr. Alopecuroid, spicâ max. Ind. Orient. Gr. Alopecuroid Indic. max. Ray Hist. est panici genus.

5. Gr. parvum repens. δυσαχυόφορον. Jamaicense spicâ longiori compreſſa.

6. Gr. Μυλοικόφορον ὀξύφυλλον. Carolinianum, s. Gr. altiſs. panic. maximâ speciosa, e spicis majorib; compreſſiuſculis utrinq. pinnatis Blattam molendinariam, quodammodò referentibus, compositâ; folijs convolutis, mucrone, pungentib; an Gr. marit. max. Curaſſavic. paniculis divisis P. B. P. hujus ad exsculpend. nobile specimen nobis dedit Insignis & in omni Naturali scientiâ vir, magnâ judicij soliditate, si quis alius præcellens D. Gulielm; Courtene, olim condiscipulus, & semper mei amiciſſimus.

7. Gr. exile duriusculū marit. foliolis circumvolutis veluti junceis brevibus. Gr. pumilum Loliaceo simile, Ray Synop. Stirp. Brit.



Tab. xxxiii.

f. 1. Gramen spica Brizæ minus, C B P. I B. Park.

2. Gr. hordeaceum à maritimis pumilum. Gr. secalinum palustre, & maritim. Ray Hist. an Gr. Hordeo disticho simile C B P.

3. Gr. caninum marit. spicatum C B P. Gr. can. mar. spicat. Monspeliens. Park. Gr. can. marit. spicat. Casparo Bauhino I B.

4. Gr. marit. spica loliacea fol. pungentib; nostras an Gr. spartaum spicat. fol. mucronatis; vel spica Secalina C B P. Spartum spicatum pungens Oceanicum I B.

5. Gr. Phalaroid. spicâ brevi reclinatâ ex utriculis prodeunte, an Gr. Phalaroid Lusitanic. Ray Hist.

6. Gramen Alopecuro; accedens ex culmi geniculis, spicas cum petiolis longiusculis promens. Gr. serotinum arvense spicâ laxâ pyramidali, Ray Hist.

7. Gr. mont. hirsut. folijs. spicâ leucophæâ dirupta. Gr. pumilum hirsut. spicâ purpuro-argentea molli nostras Ray Hist. His duab; noviſsimē memoratis Graminum specieb; multisannis nos ditavit ex propria collectione amiciſsimus vir D. Samuel Doody. Pharmaceutices & Rei Herbariæ callentiſsimus.

8. Gr. typhinum marit. minus Ray Hist. an Gr. typhin. Danicum minus Park. quærit & ipse Clariſs. Rajus.

9. Gr. parvum præcox spicâ laxâ canescente Ray Cat. Pl. Angl.

10. Gr. paniculat. bromoid. minus, paniculis aristat. unam partem, spectantib; Ray Hist.



Tab. XXXIV.

1. Gr. avenaceum parvum procumb. paniculis non aristatis Ray Hist. Pl. univers.
2. Gr. capillaceum locustellis pennatis non aristatis, Ray Hist. Gr. junceum Dalechampii Hist. Lugd. Gr. foliolis junceis brevibus radice nigrâ & albâ C.B. pin & prodr.
3. Gr. cyperoid. angustif. spicis parvis ſeſſilib, in foliorum alis Ray Hist.
4. Gr. cyp. elegans spicâ compositâ, Ray Cat. Pl. Ang. ubi ad spicam (dicit) nullum apponitur folium, & in Hist. (spicâ compositâ asperiore) legitur, = nostri tamen spica mollis est, & foliolum ei subjectum habet.
5. Gr. cyp. polystachion flavicans spicis brevib, prope summitatem caulis Ray Hist.
6. Gr. cyp. polystachion lanuginosum Ray Hist. an Gr. cyperoid. Norvegicum parum lanosum, Park.
7. Gramini cyperoidi ex monte Ballon simile, spicâ total. e. plurimis spicis compositâ Ray Hist. an Gr. cyp. elegans multiferâ spicâ Park.
8. Gramen cyper. ex monte Ballon simile humilius, in maritimis & arenosis naſcens, Ray Hist.
9. Gr. cyperoides spicâ simplici compreſſâ distichâ D. Plukenet. Ray Synop. Stirp. Brit. & Hist.
10. Gr. cyp. minimum sentinib, deorsum reflexis, pulicifornibus. Ray Hist.
11. Gramini Luzulæ accedens glabrum in palustribus proveniens paniculatum, an Gr. cyp. palustre leucanthemum Ray Hist.



Tab. xxxv.

f. 1. Gr. junceum clavatum minim. s. Holoſteum paluſtre repens, folijs, capitulis, & ſeminib; Psyllij Ray. Synops. Stirp. Brit. & Hist. à Reverendo viro D. Matthæo Dodsworth, Evangelij prædicatore vigilantiſsimo, & Botanicorum numero nequaquàm poſtremo, nobis primùm oſtenditur.

2. Gram. junceum. s. Holoſteum minimum paluſtre capitulis quatuor longiſsim. staminib; donatis Ray. Syn. Stirp. Br. & Hist.

3. Guaiacum Jamaicense Lentiſci subrotundis folijs læte virentib; flore albo. Guaicū Officinar. Parad. Bat. Prod. Guaiac. folijs Lentiſci = C B P. Guaiaci ramulus Clus. ad Monardem. Guaiac. Terentij apud Hernāndez.

4. Guaicū Jamaic. folijs veluti muriâ conditis, ſpiſsiùs virentib; flore subcæruleo. an Guaiacum folijs Ceratoniæ, Aceris. s. Burſæ paſtoris fructu, flore cæruleo Breyn. Prodr. 2.

5. Gratiolæ affinis è Coromandel folijs ſuperiorib; Rubiæ quadrifoliæ crenatis, imis, Coriandri modo tenuiſsimē diviſis.



Tab. XXXVI.

1. *Halimus humilis erecta ſemine folliculis membranaceis, bivalvib, in latitudinem porrectis, & utrinq recurvis, longo pedunculo inſidentib; clauso. Hal. anguſtif. ſeminib; ſingularib; vasculis Burſæ paſtoriæ æmulis bicornib; incluſis. Ray. Hiſt. app. an Halim. min. Germanic; Park Halim; min. CB. pin. et Halim; aurei arvi ejuſd. prodr. hanc etiam in maritimis Belgiis nuperrime invenit Spectatiſsim. vir. et Rei Herbariæ amantiſsimus D. Carol; du Bois cuj; ramuli ſummitas ſatis infeliciter exſculpta apud Ioh. Bauhin. ad Portulacæ marinæ fruticoſæ Iconem appingi videtur. itemq apud Bodæum à Stapel. Annot. in Theophr. pag. 509.*
2. *Hedera monophylla Convolvuli ſolijs virginiana. Clematis hederacea virginienſis umbilicato ſolio flore papposo Cat. HRP. Hedera virginian. Clematis facie, radice flaveſcente, Cat. HAL. ſi huj, flores pappis coronantur ad Clematitis genus amandari debet multoties hanc plantam nos florentem vidim; at nuſquam nobis Pappus occurrit.*
3. *Heliotrop. monoſperm. Indic. procumbens, glaucophyllon florib; albis. an Totecy-Xochiuh Mexic. Hernand. ? Heliotrop. Americ. minus glabrum folio, anguſto glauco. Br. Pr. 2.*
4. *Hemionitis American. ſpecioſa ſolio Securis Romanæ figuram æmulante. Hemionitis parva Antegoana ſolio Platani ferè in modum diviſo Ray Hiſt. app. ab honeſtiſsimo viro D. G. Horsnel ex Antego Inſula in Patriam reduce nos accepimus.*
5. *Hemionit. Ind. Orient. Phagi s. Eſculi ſolio aliquatenus accedens. an Hemionitis multifida Zeylan. Coſtâ ſagittatâ s. Filix ſolijs integris Zeylanica pediculo alato. Breyn Prod. 2.*
6. *Hemionitis pumila ſolijs integris, trifidis, et quinqfidis varia Cambrobritannica. Hemionit. pumila trifolia vel quinq ſolia maritima D. Lloyd. Ray Synop. Stirp. Brit.*
7. *Hemionitidi affinis Americana epiphyllanthos, ſolio ſimpliciter pinnato, Hippogloſsi æmulo, radice repatrice lignoſâ, ad foliorum crenas florida.*



1. Hesperis prodigiosa hortensis Molluginis ad imaginem degenerans.
2. Hieracium folijs & floribus Dentis Leonis bulbosi semine curvo. Moris. Prælud. Dens Leon. folijs radialis Bot. Monsp. Dens Leon. minor folijs radiatis. CB. prodr. & pin. cui & Trinciatella Italorum Camerar. Hort. Med.
3. Hieracium s. Pilosellæ majoris species folijs rariūs dentatis flore singulari nostras.
4. Holosteum s. Coronopus serpentina folijs betonicæ angustis, spicâ Rubiæ spicatæ gracili. an Holost. strictiſsimo folio minus. C B P. serpentina ovium minim. Lob. Icon. Holosteum nobis hic allatum optimè convenit cum Descriptione Plantæ quam Ioh. Bauhin. in Hist. pro Coronopo serpentaria ovium minima proposuit in quâ tamen foliola cauliculis adnexa non concedet.
5. Hyosciamus pusillus aureus American; Antirrhini folijs glabris. fortè Hyosciami albi Ægyptij. Veslingiani. varietas minor ex Americâ. hujus unam aut alteram plantam à seminib, ex Americâ delatis curâ & vigilantiâ D. Georgij London. Regij nunc Hortulani, & de Rebus Botanicis optimè meriti, pronatam conspexim; in Horto cultiſsimo novisq; ac rarioribus undiq; conquisitis stirpibus instructiſsimo, Honoratiſsimi Domini, & Reverendi admodum in Christo Patris Henrici Compton. Episcopi Londinensis, omnium Botanophilorum Mecœnatis colendiſsimi.



Tab. XXXVIII.

1. *Jacea latifol. caule alato capite magno turbinato, multiplici aculeo deflexo è squamularum cuspidib; orto. Jacea latifolia spinos. capitulis pulchris flore purpureo* I.B. *an Jac. Cyanoid, altera alato caule. Cyanus Creticus spinos; Belgarum,* Cat. H.R.P. *Jacea folijs Seridis candicantib; purpur.* C.B.P. *Jacea purpurea valentina muricato capite Clus. cui & ille attribuit folia Seridis candicantia.*
2. *Jacea montan. alato caule purpurea ex singulis capitulor. squamis spinulis. plurimis in Hemicyclum dispositis. Jacea mont. purp. echinato capite* Park. *Jacea Cyanoid. echinato capite,* C.B.P. *Jac. mont. echin. capite Clus. forte etiam Jacea capitulis hirsut. Boelij Ger. emac. sub hoc enim titulo multis abhinc annis et à D.* Edvard. Morgan. *viro in Rebus Botanicis haud infimæ notæ accepimus.*
3. *Jacea spinosa Sonchi augusto folio, maritima, minor, ad squamas capituli, spinulis plurimis.*
4. *Jac. incana Cyani majoris folio in aculeum desinente, squammis cum ciliarib; nigris. an Jac. nigr. angustif, vel Lithospermi arvens. fol.* C.B.P. *quæ 1.ª in prodr. item Jac. nigra angustif. alt. fol. longiorib; lanuginosis. ejusd. prodr. forte etiam Jac. min. fol. aculeatis, Cyani majoris nomine missa ejusd. Pin.*
5. *Jac. Tingitana Centauroid. echinato capite villoso, ex unaquaq. squama plurimis aculeis circulariter radiatis. Huic proximè accedit & ni fallor eadem est Jac. capite longis aculeis spinoso* C.B.P. *Quam iste vir magnus & in arte Botanicâ δ Ἰσιϭμέγισθ, à Jac. mont. echinato capite Clus. diversam esse, rectè pronunciabat. Centaurij marini nomine ad Casparū. Bauhinum D. Doldius misit.*

TAB. XXXIX.



Tab. xxx. ix.

1. Jac. laciniato Sonchi fol. s. latifolia purp. capitulo spinos. C B P. ex capitulor. squamis spinulæ plures pulchrô ordine stellam dimidiatam referentes, atq; deorsùm tendentes exeunt, quarum media cæteris latior & longior evadit.
2. Jac. min. Stoebe Gallicæ & Austriacæ accedens, squamis capitulor. crebrioribus, quarū singulæ unico saltem aculeo sunt armatæ.
3. Jac. min. exotica ad foliorum margines spinulis donata.
4. Jac. min. Attica capitulis plurimis ex apicibus squamularum spinâ rectâ singulari. ā doctiſsimi pariter ac humaniſs. necnon Reverendi admodum viri D. Georgy Wheeler. Eq. aur. liberalitate accepim.
5. Jacobæa lanosa folijs brevibus subrotundis Pilosellæ flore impense flavo. an Jacobæa lanata folijs brevibus subrotund. D. Ban. Cat. Stirp. Virg.
6. Jacob. pumila Absinthij Umbelliferi folijs incanis, florib; velut in Umbellam positis. an Chrysanthemum Alpin. 1.m Clus.
7. Jacob. pumila sinuatis folijs argentea, summo caule flore singulari. Ingenioso viro & Chirurgo Londinensi D. Pareo hujus communicationem debeo.
8. Iberis Americana latiorib; serratis folijs, ad caulem virticillatis.



Tab. XL.

1. Juncus acutus maritim. caule triquetro rigido, mucrone pungente, an Juncus acutus maritimus caule triangulo. CBP. 11. & Prodr. 22.

2. Junc. acutus maritim. caule triquetro maximo molli, & procerior nostras.

3. Junc. acutus maritim. caule triquetro molli, spicâ glomeratâ in Stylo ſeſſili.

4. Juncus acutus maritimus capitulis rotundis C B P. 11. Juncus acutus maritim. alter Park.


5. Juncus minor acutus maritimus prolifer. s. ex uno capitulo multa.

6. Juncus parvus paluſtris cum parvis capitulis Equiseti, Ray Hist. Junc. mont. parvus cum parvis capitulis luteis I B. Gramen sparteum capitulis Equiseti. Merett. Pin. Rer. Nat. Brit.

7. Juncelli ovum minimi capitulis Equiset Ray Hist. primus ejus Inventor fuit amiciſsimus vir Reiq. Herbariæ amantiſsimus D. Matth. Dodsworth, qui et jam olim hujus exemplar, optimè exsiccatum, nobis benignè communicavit.

8. Junc. parvus calamo supra paniculam compactam longiùs producta Raij Hist. in Westmorlandiâ invenit ille Stirp. Britannicar. explorator indefeſsus, D. Iacob. Newton. cujus munere accipiebam.

9. Juniperus Caroliniana Thuyæ ramulis fusis, & Compreſsis, odoratior. an Cedrus folio Cupreſsi major, fructu flavescente CBP. Thuyæ genus 4^m. Hist Lugd. 61.



Tab XLI.

f. 1. *Lamium album Parietariæ folio virginianum. Lamium Parietariæ facie* Moris. Præl. Bot. 278. *Lamii Parietariæ folijs* Ic. Roberti. *sed hujus Icon minus placet. an Lamium album non fœtens folio ablongo minus* Cat. HRP. 99. *Lamium Parietariæ folio* Bruynero Lonquet Hort. 69.

2. *Lamium pusillum rubrum Paridariæ facie Americanum. an Lamium Novæ Angliæ Parietariæ folijs* Raij Hist. 560.

3. *Lam. rubr. minus folijs profunde incisis* Raij Hist 560.

4. *Lam. Cannabinum aculeatum flore speciosa luteo, labijs purpureis. Cannab. spuria flor. speciosa luteo, labro purpureo* Ray Hist. 561. *an Cannabis altera spuria* Park. pag. 599. *an Urtica aculeata folijs serratis altera* CBP. 232. *quærit & ipse* Cl. Rajus.

5. *Lappula Bengalens. tetraspermos Ribesij folio echinis orbicularib; ad folior exortus plurimis simul Seſsilibus. an Alcea Indiæ Orientalis fructu lappaceo folio Ribes* Breyn. Pr. *Nexpayan Mexicensuim* Hernand *huc pertinere videtur.*

6. *Lenticula palustris Indica foliolis subrotundis binis capillamentis ad imum barbatis.*

TAB. XLII.



f. 1. Leucoium. s. Lunaria basculo sublongo intorto. Raÿ Hist. 789.
2. Lichen parvus erectus foliolis profundè laciniatis. an, Lichen minim. foliolis laciniatis. Raÿ Hist. 126. in Umbrosis humidis ad parvulum fluentum juxta Fontes medicatos apud Dovvicum, quatuor circiter milliaria à Londino. diſſitum, in agro Surrienſi, copioſiſſimum invenimus.
3. Lilio-narciſsus, s. Narciſsus Liliflorus Carolinianus. flore albo, singulari, cum rubedine diluto. ab honestiſs. viro & Hortulano admodum industrio. D. Gulielm Darbÿ. recepimus ad quem huj. radices plurimæ ab amico suo è Carolinâ allatæ sunt, & in viridario suo jam nunc superbiant apud, Hogseden, vicum Londini, ad aquilonem suburbanum. lit. a. capsulam triquetram seminib; oppletam significat.
4. Limonium reticulatum supinum Boccon. de Pl. Sicul. pag. 82.
5. Limonium minimum comatum elegans. an Limon minus densâ caulium feniculaceâ veluti comâ Lœl. Triumfet. catal. apud Fratrem. p. 65. fortè etiam Limon. Ferulæ folio. Boelÿ. Park. 1235.
6. Liquid-ambari arbor. s. Styracifera Aceris folio fructu tribuloide, (i e) pericarpio orbiculari ex quamplurimis apicibus coagmentato semen recondens. Ocosol. Indorum Monard. apud Clus. 302 Xochicotzo Quahuitl Mexicens. Hernand. Styrax Aceris folio Hon. Belli. CB. prodr. Styrax arbor virginiana Aceris fol. Raÿ Hist. nec Acer nec Platanus propriè dici potest; est enim quædam Heteroclyta & sui generis Arbor. lit. a. fructum muricatum; cujus et transversam sectionem. b. indicat c c c. semina.
7. Lithospermum angustifol. umbellatum CBP. 258 & in append. 521. Park. Paul. Boccon. de Pl. Sicul. pag. 77.
8. Lychnis Alpina muscosis folijs densiùs stipatis, floribus parvis, Calyce duriore ab Ocymoide muscoso Park. plurimum differt.



Tab. XLIII.

f. 1. *Lotus argentea Cretica. Lotus πολυκέρατος fruticosa Cretica argentea, siliquis longiſs. propendentib; rectis* Moris. Hist. 2. *(nostræ tamen siliquæ pendulæ non sunt sed sursum rigescunt.) an Lotus incanus major s. Trifol. Corniculatum siliquâ grandi, incanum maritim.* Raÿ Hist.

2. *Lotus fruticosior Africana, folijs incanis, floribus binis, amplis, coccineis. à D. Ioanne* Fishwick *accepimus.*

3. *Lotus tetralobos minor Creticus. s. Lotus parva, Cretica, procumbens, incana, quaternis siliquis cruciatim dispositis. Candie Lotocruce.*

4. *Lychnis Caryophyllæus virginian; Gentianæ folijs glabris, quatuor ex singulis geniculis caulem amplexantibus, flore amplo fimbriato. noster.* Raÿ Hist. app. 1895. *huc fortè pertinet Eloquiltic Mexicensibus. s. Herba geniculata* Hernand. 210.

5. *Lysimachiæ Ephemeron dictæ, genus Maderaspatense, Limonij folijs subrotundis, caule craſso fungoso. lit. a. florē expansū ostendit.*

6. *Lysimachiæ spicatæ purpureæ affinis Thymifolia, flosculis in cacumine plurimis, quaſi in nodis junctis. Anthyllis fortè* Prosp. Alpin. *exot. hujus specimen Reverend. vir D. Geo. Wheeler. è Græciâ (quâ invenit) rediens nobis munere donavit.*

7. *Tithymalus botryoid. Maderaspatens Hyperici folijs non crenatis, florib; ex alis uno versu provenientibus inter Lysimachias in Tabulis nostris per errorem transpositus.*

TAB. XLIV.



Tab. XLIIII.

f. 1. *Lyſimachia non papposa Meliſſophylla Maderaspatens. forte Melochiæ genus ſiliquâ quinq; capsulari.*

2. *Malva Tingitana. flore Cæruleo parva.*

3. *Malv. rotundifolia. glabra ex Hispaniâ flore amplo rubello.*

4. *Malv. Gallo-provinciæ laciniata s. Alcea tenuiſſim. Pulsatillæ folijs. flor. ampl. rubente, Alcea tenuifol. humil. maritim. Gallo-provincialis fol. inferiorib; nonnihil ad Geranium accedentib; Sch. Bot.*

5. *Malv. Oriental. elatior longiori fol. fl. alb. minimo, capsulis conglomeratis. an Malv. frutesc. Melochiæ angusto fol. florib; alb. fasciculatim. congestis. Par. Bat. Prodr. inter addenda.*

6. *Mandarù Maderaspatense folijs firmiorib; parvis bisulcis, glabritie splendentib; ad Surculum densiùs stipatis.*

7. *Muscus polytrichoides elatior folijs angustis pellucidis & ferè membranaceis noster* Ray. Synops. Stirp. Brit.



Tab XLV.

1. *Marrubium aquaticum s. aquatile foliis tenuiùs diſſectis* Moris. Prælud. *fortè Marrub. paluſtre hirſut. tenuiùs laciniatum* CB Pin. & in Prodr. *noſtri tamen folia ſunt omnino glabra.*
2. *Marrub. an potiùs Cardiaca foliorum apicib, in prælongam barbulam extenuatis, verticillis florum petiolis innitentibus. an Marrub.* ex Aleppo *verticillis longis pediculis inſidentibus.* Cat. Hort. Edinburg.
3. *Matricariæ* Achaovan. *dictæ ſimilis, Eryſimi foliis, Abſinthii ſapore* Jamaicenſis.
4. *Melilotus lutea* Indiæ. *Orientalis erecta, folliculis rotundis, parvis, ſpicis florum ex foliorũ alis multiplicatis.*
5. *Melilot. Indic. humil. erecta florib, exiguis odorat. albis, pericarpiis majorib, ſpicatim denſius ſtipatis.* Suendadi-pullu Hort. Malab. P. 9.
6. *Mentha hirſutior* Βραχυσταχυφυλλος *exotica à D:* Doody. *accepimus*
7. *Muſcus polytrichoides humilior foliis latis ſubrotundis. Adianthum aureum humilius fol. latis ſubrot.* Raii Hist. *an Muſc. Trichomanis facie Jungermanni foliis utrinq. rotund. ſplendentib,* Cat. Aldtorf.
8. *Melilot. lutea minima procumbens foliis minoribus crenatis. an Melilot. Meſſanenſis procumb. folliculis rugoſis ſublong. ſpicis florum brevioribus,* Ray Hist. Cat. Exterar.



Tab. XLVI.


f. 1. *Meſpilus aculeata Pyrifolia denticulata ſplendens fructu insigni rutilo Virginienſis. an Meſpil; virginiana colore rutilo CBP. hujus folia aliquando variant, & senectute multùm laciniata evadunt, ut in Horto Nobiliſs. viri D. Caroli Howard. prope oppidum Dorkin in agro Surrienſi obſervavimus. hujus folium ad lit. a. in Tab. exprimitur.*

2. *Meſpilus Prunifolia Virginiana, non ſpinosa fructu nigricante. Rhamnus virginiana Prunifolia fructu nigro, oſsiculo compreſso Banist. Cat. Stirp. Virgin. an Meſpilus, s. Sorbus torminalis virginiana folio latiſsi fructu nigro Sch. Botan. fortè etiam Meſpilus spinosa s. Oxyacantha virginiana nigra Cat. HAL. si saltèm aliquandò inveniatur spinosa, & ubi magis domestica spinis careat, ut accidit in Anthedone Azarolo Cæsalp. Theophraſto monente. nostram autem unquam spinoſam eſse, me haud perſvadere poſsum, b. ramulum fructu onustum significat.*


3. *Meſpilus Virginiana. Apij folio vulgari similis major grandioribus spinis.*

4. *Meſpil; Apij folio Virginiana spinis horrida, fructu amplo coccineo. Meſpilus Apij folio Indica Sch. Bot. Meſpil; spinos. s. Oxyacantha maxima virginiana Cat. HAL. Oxyacantha Oſsiculis binis fructu magno inclusis. Banist. Cat. Stirp. Virg. Oxyacantha Americ. Galli calcar dicta. R.S. Ejus Ray Hist. hujus Arbor Spina Glaſsenburienſis, Spina Christi, & Spina Sancta per errorem dici solita est. lit. a. Surculum cum aculeo armatum b. b. oſsicula demonstrant.*

TAB. XLVII.



Tab. XLVII.
f. 1. Millefolium Tanaceti foliis flore albo Moris. Præluď. an Corymbifera Millefolij Umbellâ, folio alato & laciniata Raӱ Hist.
2. Muscus clavatus Ceranoides, segmentis latioribus, Compreſsis, eburneo nitens, noster Raӱ. Syn. Stirp. Brit. app.
3. Musc. clavatus Ceranoides Gagateus nigredine splendens, noster Raӱ. Syn. St. Brit. append.
4. Musc, montanus, gracilis, ramosus, bifidulis longioribus glabris. an Muscus ramosus erectus minor CB. pin. & prod.
5. Musc, Alpin, ramosior, erectus, flagellis brevioribus, lanuginosus. an Muscus terrestris ramosus erectus major CB. pin. & prodr. Muscus montanus Tab. Icon. fortè etiam Muscus hirsutus capillaceus Merett. Pinac.
6. Muscus parvus stellaris CBP. Park. Adianthi capitulis. Muscus in ericetis proveniens IB. Hist. Lob. Icon. sed figura non placet.
7. Muscus terrestris erectus minor, polyspermos Raӱ Hist. & Synops. Method. Stirp. Brit.
8. Musc, terrestris repens pediculis foliaceis binis clavis in altum se erigentibus. Muscus alius terrestris, Lycopodium Dodonæi Hist. Lugd. qui à Musco terrestri 2. ejusd. Histor. (ie) terrestri repente à Trago picto IB. plurimum differt, licet in Pinace suo Casp. Bauhin, ab illo non separat, sed planè ut unum confundit.
9. Muscus Zeylanic, erectus, perpetuò virens, in arboris proceritatem excrescens. an Quaniazatl. s. Spica arboris Hernand. Hist. Mexic.
10. Muscus Coralloides, multifido capillaceo folio, Palmites pelagica nigra. Musc, capillac. multifid, niger. CBP. adlit. a. Palmites pelagica alba depingitur. Muscus capillac. multifido folio albido. CBP. Corallinæ affinis s. Muscus marinus tenui capillo. IB. Muscus marinus bulgatissim, Lob. Icon. p. 2.
11. Muscus marinus denticulatus, procumbens, caule tenuiſsimo, denticulis bijugis Raӱ. Histor.
12. Musc, pennatus, ramulis & capillamentis falcatis Raӱ Hist.
13. Muscus marinus Neapolitanus Park. Coralloides lenta foeniculacea IB. Fucus foeniculaceus colore livido nobis Raij Synop. app.



Tab. XLVIII.

f. 1. Muscus aureus, capillaris, paluſtris, inter foliola folliculis rotundis (ex ſententiâ D. Doodij) quadripartitis. Graminifolia paluſtris repens, baſculis granorum Piperis æmulis. Raÿ. Hist. & Synops.

2. Muscus pelagic; pennat; rubens, ramulis numeroſis mollib; latiùs ſe ſpargentibus. an Musc; marin; rubens, pennatus noſtras. Raÿ Hiſt. item Musc; marin; purpur. parvus, folijs oblongis. Millefolij ferè diviſurâ. ejusd. Hiſt.

3. Musc; marin; denticulatus minor, ramulis in creberrima capillamenta ſparſis. an Muscus marinus denticulat; minor, denticulis alternis Raÿ Hiſt.

4. Muscus marinus geniculatus, eburneo nitore candicans, erectus, cauliculis pennatis. Muscus marinus candidus Coralloides Hiſt. Lugd. Lob. Icon. p. 2. Corallina altera Tabern. Icon. ſed figura minus placet. Muscus Coralloid. ſquammulis loricatus. CBP.

5. Muscus marinus major argutè denticulatus Raÿ. Hiſt. Muscus Coralloides alter albus Lob. Icon. part. 2. Abies marina Belgica Clus. Hiſt. Abieti ſimilis maritima CBP.

6. Muscus marin. s. Coralloid. non ramoſus erectus, arenacei coloris, Aſtacorum corniculi adinſtar. geniculatus. Muscus marinus Equiſeti formis non ramoſus Raÿ. Hiſt.

7. Myoſotis Scorpioides. an potius Heliotropium minus erectum, folijs ſubtus incanis, è Maderaſpatan.

8. Myrt; Brabanticæ accedens Africana, baccis carens, Conifera. Arbor Conifera odorata folijs Salicis, rigidis, leviter ſerratis Raÿ. Hiſt. an Laur; ſerrata odora Promont. Bon. Spei. Bod à Stapel. in Theophr. p. 333. ex America quoq; nobis allata eſt, & à Noſtratib; Inſulam Bermudenſem vigentibus, Laurus odorata, vulgò nominatur, ut ab honeſtiſſ. viro Jacobo Harlow. didicimus.

9. Myrt; Brabanticæ ſimilis. Caroliniensis, baccifera, fructu racemoſo, ſeſſili, monopyrene, fructum fert veluti Saccharo candidiſſ. incruſtatum, cujus ramulus cum fructu, ut per Microſcopium apparuit, adpingitur. Hujus pulchri Sanè, perrari, & prorſus novi Arbusculi Americani generis, munificentiâ Illuſtris viri D. Gulielmi Courtene, compos factus ſum.



Tab. XLIX.


f. 1. *Muscus polytrichoides anguſtifolius pellucidus ramoſus noſter* Ray. Synops. *Stirp*. Brit.

2. *Njara Hort. Malab. ſimilis è Maderaspatàn. Arbor exot. Ceratiæ fol. Corymbosa.*

3. *Nir-Schulli Hort. Malab. accedens è Cormandel. Gratiolæ affinis Indiæ Orientalis, Digitalis æmula. ubi tres exhibentur ſpecies, ſeminum figurâ diſcrepantes, ut literæ a. b. ccc. denotant.*

4. *Oenanthe aquatica, Pimpinellæ Saxifragæ diviſurâ, noſtras. Oenanthe Staphylini folio aliquatènus accedens* IB. *Oenanthe Paſtinacæ Sylveſtr. folio ſemine Atriplicis* CBP.

5. *Onobrychis Biſnagarica, Mimoſæ folijs, ſiliquis ad unum ſolummodò latus dentatis, ex alis polyceratos.*



Tab. L.

f. 1. *Onobrychis minor, flore parvo, purpureo, siliquâ echinatâ, majoribus aculeis.* Ray Hist. *Onobrych. minor, siliculis echinatis, cristatis, majorib; & craſsiorib; aculeis prœditis donata* Moris. Hist. 2. *an Onobrych. fructu echinato minor.* CBP. *item Caput Gallinaceum minus ejusd. pin. & prodr. Onobrychis. s. Caput gallinaceum, minus, fructu maximo, insigniter echinato.* Lely. *Triumfelli. Cat. apud fratrem.*

2. *Onobrychis minor foliolis cordatis, siliquis magnis, asperis, compreſsis, beluti isthmo interceptis, floribus in globum congestis. an Onobrychis folijs viciæ, fructu echinato major caulescens, & ramosa* Schol. Botan. *an Onobrychis semine Clypeato aspero minor* CB. Pin. & Prodr.

3. *Onobrychis Gangetica monophyllos, siliculis singularibus, lævibus, falcatim, per internodia discriminalis.*

4. *Pai-paroea, s. Couradi* Hort. Malab. *similis, sinon eadem. Frutex baccifer Malabaric; fructu plano rotundo piloso tetrapyreno.* Ray Hist.

5. *Perfoliata alpina angustifolia minor* CBP. *quœ 4.ª in* Prodromo. Park. Theatr.

6. *Perfoliata minor angustifolia, Bupleuri folio* CBP. *Perfoliatum angustifolium montanum* Col. ecphr. 247. *Perfoliata angustifolia Alpina minima* Park. *Auriculæ leporis affinis, Odontitis lutea* Valerandi. & Dalechampy. IB. Hist. tom. 3. lib. 27. p. 201.



Tab. LI.

f. 1. *Palma Jamaicens. non spinoſa humilis, folio multiſciſſo, flabelli formi, Ampana. Malabar. haud abſimili*

2. *Phaseolus maritimus purgans, radice bivaci, folijs craſsis, subrotundis, Bisnagaric, an Phaseolus Braſilicus Florileg. Renovat. edit. p. Matth. Meriani. Katu-tsjandi. H. Malab. p. 8.*

3. *Phaseolus Orientalis Anagyridis folio, ſiliquâ falcatâ glabrâ.*

4. *Phaseol; minimus Maderaspatan; florib; miniatis Meliloti adinstar spicatis, huc fortè pertinet Ystactlalamatl Mexicens. Hernand. Hist. 452.*

5. *Paronychia myosotis virginiana, folijs subrotundis. Nasturtium myosotis insipidum Banist. Cat. Stirp. Virgin. à D. Bobert. amiciſſimè communicatum fuit, hujus optimè aſervatum exemplar.*



Tab. LII.
f. 1. Ornithopodium majus è Maderaspatàn. folijs lanugine argentatis.
2. Phaseolus parvus spicatus Maderaspatens. cordatis, & incanis folijs, siliquâ utriculatâ nigrâ.
3. Phaseol; minor Vinagericus folijs argenteo colore villosis, siliquis torosis brevibus, spadiceâ hirsutie pubescent. fructu parvo Scaraboide nigro. an Phaseol; Zeylan. Salviæ folijs subrotund. mollib; & hirsutis Parad. Bat. Prodr. item Orob; Phaseolôdes Zeylan. folijs Salviæ rotundis incanis subhirsutis ejusd. P.B.P.
4. Phaseolus Nevisensis. folijs multangulis, tuberosâ radice esculentâ, Phaseolus perennis Martinianus radice tuberosâ esculentâ, anguloso crenato folio P.B.P. nostratib; the Turneptree vocatur.
5. Phaseolis accedens, Frutex Iaphnapatensis spinosus, fructu Ovali, nigris & cinereis punctis, elegantèr guttato P.B.P.
6. Pilosella minima, Gnaphalij montani lanosis folijs, flore aureo, nondùm descripta.
7. Polygonum minus, procumbens, niveum, facie Glauris exiguæ, Polygonum minus candicans supinum Botan. Monspel. Polygala repens nivea C.B.P. Polygala repens Nuperorum Obs. & Ic. Lob.



Tab. LIII.

f. 1. Padri Malabaricæ similis, Arbor è Moderaspatan. (h. e) Clematis arborea Indica Iuglandis folio, flore luteo, spicato, minimo, siliquis longis, angustis, quadratis, atq; contortis, seminibus substantiâ lignosâ obductis Breyn. Prodr. 2.

2. Polygalæ affinis Africana glabra, floris galeâ binis cristis fimbriatis, ornata, ex involucro magno dipetalo erumpente.

3. Polygonum maritimum longius radicatum nostras, Serpylli folio circinato, craſso, nitente, Synops. Stirp. Brit. app. fortè Polygon. minus lentifolium C. B. Pr. prodr. Polygon. minus Monspeliens. Park. huic proximè conveniret Polygon. Herniariæ folijs, & facie, per amplâ radice Astragaliti Adv. Lob. si hæc herbula non eſset tota incana, nostra autem glabritie splendet.

4. Polygonum lignosum à maritimis Africanum, azurei coloris foliorum bice, squammulis pellucidis ornatum. Limonium lignosum Gallas ferens Bocconi, huc pertinet, & fortè idem est. à D. Fiſhwick. nobis tranſmiſsū.

5. Ptarmica virginiana Scabiosæ Austriacæ folijs indiſsectis. Ptarmica virginiana folio Helenij Moris. Prælud. Pseudo costus virginiana, seu Anonymos Corimbifera virginiana flore albo Ray Hist. hujus specimen jam antè triginta circitèr annos in Horto D. Walker. apud Westmonasterium legimus, cujus filius hanc plantam cum alijs rariorib; è virginiâ transmisit.



Tab. LIIII.
F. 1. Poutaletsjæ Malabaricæ similis Arbuscula Maderaspatensis (i e) Baccifera Indica baccis oblongis in Umbellæ formam dispositis Raÿ Hist.
2. Pulegium virginianū angustifol (Serpentaria dictum) flosculis in Cymis, an potius Clinopodium Pulegij angusto rigidoq; folio virginian. flosculis in Cymis. Pulegium virginianum Nonnullis, alijs Serpentaria aut Colubrina virginiana Raÿ Hist. Satureia floribus in summitate dispositis Cat. H.A.L. Pulegio Cervino quodammodò accedens capitata Chamæpityos Austriacæ cæruleæ folijs & facie Breyn. Prodr. 2.
3. Quercus virginiana Castaneæ folio, nostra Raÿ Hist. append.
4. Quercus Esculi divisurâ, folijs amplioribus aculeatis, an Quercus alba virginiana Park.
5. Quercus virginiana rubris venis muricata The Scarlet Oake nostratib; dicta. in Horto Comptoniano, hujus plurimas, adolescentes Arbores observavimus.



Tab. LV.
f. 1 *Ranunculus hirsutus annuus flore minimo* Raÿ Hist. & Synops. Method. Stirp. Brit.
2. *Ranunc. aqualis, albus circinatis tenuiſsime diviſis folijs, florib; ex alis, longis pediculis innixis Nobis* Raÿ Synops. *Millefolium aquatium cornutum* quoad figuram I B. t. 3. *iten quoad figuram, ut & descriptionem* CB. Prod. *Millefolium aquaticum cornutum majus* Park.
3. *Rapunculus galeatus Salicis glabro folio Bisnagarius. floribus aurantijs.*
4. *Reseda Maſsiliensis folijs latioribus crispis. Reseda crispa Gallica* Boccon. 77. *Reseda latifolia flore flavo* Moris. Prælud. Botan.
5. *Reseda minor Indiæ Orient. Ptarmicæ pratensis folio, crenis exasperato, s. Sesamoides parvum Bengalense serratum.*
6. *Rhamnus Lycij Myrsinites facie Gangeticus, folijs densis, ferè convolutis, Jasmini floribus, & spinis gemellis.*
7. *Rhamnus Orientalis spinis uncinatis, atro purpureis. Oleæ. s. Ligustri folio, cujus ad imum, petiolus umbilico inarticulatur, Rhamn; Oriental. Ligustri folio spinis atro-purpureis noster* Raÿ Hist. *an Mahaleb genuinum Arabum?*



Tab LVI.

f. 1. Rhoi Obsoniorum similis, Americana, Gummi candidum fundens, non serrata, foliorum rachi medio alata. Rhus virginianum Lentisci folijs Ray Hist Banist. Cat. Stirp. Virg. Mſs. hoc est e Copalliferis Arbor, Errandi ansam nonnullis scriptorib; hæc fortè præbuit Arbor, quum Lentiscum nostrum Europæam Virginiæ plagis indigenam eſse contendere velint Vid. Hond. Geograph.

2. Ricinus Indicus pilosus. trifidus, vel quinquifid; flosculis atro purpureis, in Umbellâ aggregatis. Hoitzpochotl. Huaxacensis, s. Ricinus Novæ Hispaniæ Hernandez. Hist. Mexic. an Ricinus Americanus folio Staphisagriæ. C.B. Pin. app.

3. Euphrasia Alsines angustiori folio. Rubiæ modo Spicata, Golcondea, herbulam hanc exoticam sub Euphrasiarum vexillo militari, ab eximio Botanico D. Doody. didicimus, ideòq, malè hoc loco collocatur. ad Euphrasias ablegandа sit.

4. Saamouna Pisonis. s. Siliquosa Braſilienſis arbor, digitatis folijs, serratis, floribus Teucrij purpureis. Arbor pentaphillos virginiana floribus spicatis mono=petalis Mus. Courtenian. Ray Hist. hujus exemplar, cum multis alijs exoticis, ab insigni viro D. Gulielmo Courtene, dono accepimus.

5. Salix longiori folio hirsuta, rosea, nostras.



Tab. LVII.

f. 1. *Salvia peranguſtis ſinuatis folijs, longis pediculis inſidentibus, lanuginoſa. an Salvia anguſtifolia serrata C.B.P. Salvia Cretica anguſtifolia aurita Park. (ie) Salvia tenuifolia I.B. Hiſt. t. 3.*

2. *Salvia Cretica pomifera Clus. Hiſt. Salvia baccifera C.B.P. Salvia gallifera Cretica I.B. Hiſt. t. 3 Salvia major Cretica latifolia & anguſtifolia, aurita & non aurita, pomifera & non pomifera Park. Theatr. huic folia ſunt viridia.*

3. *Salvia auriculata mucronatis folijs craſſis, lanuginoſa, appendicibus aucta.*

4. *Salvia trifoliata minor Indica.*

5. *Salvia Ciſti folio e Promontorio Bon. Spei. forte Salvia Syriaca Ciſti famin. folijs acetabulis Moluccæ Breyn. hujus etiam Leonard. Rawolfius meminit in ſuo Hodœporico.*

6. *Salvia fruticoſa Ciſti folio haud incano, florib. purpureis. Verbaſcum verum flore rubro arboreſcens Cat. Hort. Groningens. Verbaſc. sylv. Salvifolium exoticum folio rotundiore Lobelij I.B. Hiſt. t. 3 Verbaſcum subrotundo Salviæ folio C.B.P. Verbaſc. Salvifol. exoticum Ciſtoides Park. et sub Verbaſci Ciſtoides nomine ex Horto Cl. Contareni habuit Caſpar. Bauhinus, hujus flores Lobelio videre nuſquam contigit; et Ciſti æmulos appingit Parkinſonus, item pentapetalos eſſe, petalis subrot. colore flavo, ſtaminulis rubentib. deſcribit, ſed hæc omnia per figmentum, nec minus erroneè à Cl. Hermanno in Cat. H. Ac. Lugd. append. elegans hæc & minus cognita ſpecies, Verbaſci latis Salviæ folijs C.B. Synonymo proponitur. Antè triginta circiter annos in Horto D. Morgani luxuriantem obſervavimus.*

7. *Sanicula aizoides tridactylites Alpina minor flore albo majore, Sedo Alpino tertio ſimilis, altera Fab. Col. ecphr. p. 2. Sedum Alpinum trifido folio C.B.P. Sedum Alpin. laciniatis Ajugæ folijs. Park. Sedi affinis triſulca Alpina flore albo I.B. Hiſt. t. 3.*



Tab. LVIII.

f. 1. *Sambucus humilis, s. Ebulus laciniato folio* C.B.P. *Ebulus laciniata* Ponte. *Ital.* a. *flores* b. *fructum indicat.*

2. *Sanicula myosotis alpina, florib; albicantibus, ferè Umbellatis. Saniculæ Alpinæ aliquatenus affinis* I.B. t. 3. *Cotyledon hirsuta Phyt.* Brit. Ray *Hist. et Synops. an Saniculæ montanæ alterius species* 2ª. *Clus. Hist.*

3. *Sanicula s. Cortusa Americana, spicata, floribus squallidè purpureis.*

4. *Scabiosa peregrina Thapsicæ majoris folio, squammatis capitulis, flore flavescente.*

5. *Scabiosa Bisnagarica, s. Globularia frutesc. rigidis folijs, ad radicem rotundioribus cordatis, ad caulem autèm mucrone præditis.*

6. *Scabiosa Jamajcensis hyſsopifolia s. Globularia capitulis ad geniculum, folijs plurimis, stellatim ambientibus, verticillata. fortè, Tzicatz on Tecomatl, s. Caput formicæ* Hernand. *Hist. Mexic.*

TAB. LIX.



Tab. LIX.

f. 1. Sanicula virginiensis alba folio oblongo mucronato. Saxifraga alba folio oblongo acuminato D. Banist. Cat. Mſs.


2. Schetti Hort. Malab. ē Maderaspatan. Jasminum Indicum Lauri folio inodorum. Umbellatum floribus Coccineis P.B.P. Jasmin. arboresc. Indic. flore tetrapetalo. Umbellato, phœniceo, foliis Laurinis. Breyn. Prod. 2. Ratambala Zeylanensium.

3. Scorpioides Bisnagarica folio singulari rigido siliquis articulatis erectis. Onobrychis Zeylanica folio singulari oblongo rotunda D. Hermanno. D. Sherard.

4. Scorzonera bulbosa Bistortæ minimæ foliis virginiana.

5. Scrophularia, Scorodoniæ foliis Moris. Hist. 2. & Prælud. Bot. Raij Hist. & Synops. Insectæ nostræ Angliæ. adjacentis Insulæ indigenam eſſe ā D. Sherard. nuperrime Observata.

6. Securidaca Maderaspatens. siliquis falcatis, fulvis, & villosis. plurimis circa ramulos stellatim positis. Stella Orientalis fulgens. ā. siliquam seorsum, b. semina denotant.



Tab. LX.

F. 1. Sena exigua Maderaspatana, s. Tetraphylla siliquifera glabra, florum pediculis inter foliorū exortum dispositis. an Absus Verlingÿ. Obs. in Alpin. Senna Oriental. Tetraphylla Absus dicta Cat. Hort. Ac. Lugd. Semen atro colore splendidum, parvis includit siliquis Cicerum emulis, ut ex Icone bidesis. a. semina rhomboide figurâ ostendit.

2. Sena tetraphylla, s. Absi congener hirsuta è Maderaspatan flore flavo, siliquis punctatis scabris, folliculos sub terram condens. b. folliculum c. nucleum è folliculo exutum d. siliquam cum seminibus Pyri proximis ostendunt. an Arachidna Indiæ utriusq. tetraphylla P.B.P. Mandubi Pison.

3. Senecio pusillus Maderaspatanus folijs crenatis, crassis, & hirsutis. Tlanpanxihuitl Nov. Hisp. Recch. huic haud prorsus dissimilis.

4. Seseli pratense folijs densius pilis pubescentibus.

5. Sinapistrum Indicum diphyllon, radiatis siliquis ad sinus foliorum ex adverso binis ad hujus Sinapistri genus accenseri debent. Cupa-Veela. v Aria Veela Hort. Malab. p. 9. ut nobis videtur.



Tab. LXI.

1. *Siſyrichium cæruleum parvum gladiato caule Virginian. Gladiol; cæruleus hexapetalos caule etiam gladiato Banist. Cat. Mſs.*
2. *Siſyrichium Bermudenſe florib; parvis ex cæruleo & aureo mixtis. Iris Phalangoides quorundam.*
3. *Sium minimum umbellatum folijs varijs. Sium minimum Ray Hist. Sium minimum ferulaceis folijs Moris. Prælud.*
4. *Soldæ Malabaricæ similis Arbor Maderaspatana ıulis amentis donata.*
5. *Sonchus subrotundo folio nostras, leviſsimis spinulis circa foliorum oras exasperatus. an Sonchus asper subrotundo folio C.B.P. Park. Sonchus asper vulgi, flore Hieracij Monspelienſib; Ray Hist. Monspelij sponte nascitur, & in Angliâ.*



Tab. LXII.

f. 1. *Solanum incanum Chinense minus spinosum, floribus parvis, ferè umbellatis, à D. Pettever accepi.*

2. *Solanum Indic. spinos. atro virens, & undiq; glabrum, folijs Brancæ Ursinæ divisuris, an Solan. pomiferum spinos. Indic. fructu echinato folijs Cardui s. Acanthi spinosi Breyn. Prodr. 1. in Horto quodam Hollandiæ celebriori collegit D. London, & nobis transmisit.*

3. *Solan. annuum nigricans Virginianum spinosiſs. latè se spargens, flore cæruleo glabrum. Planta spinosiſs. Virginiana Borraginis flore Tradescantio Raij Hist. 674. & in app. 1876. item Solanum pomiferum frutesc. African. spinos. nigricans Borraginis flore, folijs minus profundè laciniatis, spinis multò longioribus, majorib; & crebrioribus horridum Raij Hist. 1799. an Solan. American. laciniatum spinosiſs. P. B. P.*

4. *Sonchus ἀφυλλόκαυλος angusto & oblongo folio nostras, per foliorum ambitum creberrimis spinulis asperatus.*

5. *Sonchus angustifolius maritimus C. B. Pin. & in Prodr.*




Tab LXIII.

f. 1. *Solano àn potius Mirabili Peruvianæ affinis tinctoria Betæfolia scandens Cochin-Chinensis, flosculis puniceis muscosis spicatis, fructu intùs Cochleato. Solanum scandens Malabaric. Betæfolia Commelin. Hort. Amst. Solanum peregrin. Betæfolia* RAY Hist. *Beta baccifera aizoides rotundi folia Zeylanica Breyn. Prodr. 2. Basila Zanon. Basella Hort. Malab. P. 7. ubi sub Bryoniæ genere (minùs aptè) recenseri bellet Commelin in Not. suprà loc. Bintal Zeylanensib; & Cochin-Chinensibus Brotho-Condolo nominatur, succus ex florib; non ex fructibus (ut bult Annotator) expreſsus commodiſsimè pro miniaturà utitur. in Horto Comptoniano apud Fulhamenses cum floribus, & fructu collegimus.*

2. *Spiræa Theophrasti similis Frutex Indicus latiori folio, spicis florum è foliorum alis, hujus ramulum noster amicus Reverend. D. Dodsworth. in horto cujusdam ingeniosi biri D. Wilkinson. Pharmacopæi Eboracensis collectum & optimè reſiccatum absq; titulo nobis Londino transmisit, & fortè Evonymi nostræ Jujubinis folijs Caroliniensis barietas haberi poſsit. àn Frutex Virginianus floribus albis instar Spiræa compactis Cat. MSS. Hort. Oxon. lit. a. significat flosculos per Microscopium bisos, in perianthijs occlusos & nondùm explicatos.*

3. *Sphondylium hirsutum minus folijs tenuitèr laciniatis seminelato, Sphondyl. mont. min. angustifol. tenuitèr laciniat. nostrum* RAY. *Synops. an Sphondylium Alpinum parvum CB. pin. & prodr.*

4. *Staphylodendron Æthiopic.* Μονολαοτοκοαλλιωομονοφυλλον. *singulari hirsuto folio nitente, Staphylod. Afric. sempervirens folijs splendentib; Commel. Hort. Amst. Staphylod. Afric. folio lucido* P.B.P



Tab. LXIV.

1 . *Stachys angustifolia Lychnitis, non crenata, Stachys Cretica C. B. P. Stach. s. Pseudo Stachys Cretica Park.*

2 . *Stachys Lychnitis angustifolia, crenata, Maderaspatana, lit. a, imum folium cum ejus pediculo oblongo significat.*

3 . *Syringa Lauri-folia Jamaicensis floribus ex flavo-pallescentibus, huc fortè pertinet Totoncapalli Mexicensib; Syringa lutea Hernand. Hist. Mexic.*

4 . a. *Tamarind; Americanus siliquâ ferè orbiculatâ, monospermos Mus. Courten. Tamarind; Occidentalis siliquâ subrotundâ, Carandas fortè Bontio Ray Hist.*
b. *Tamarindus Bisnagaricus siliquâ productiore dispermos, seminib; veluti isthmo interceptis, c.c. semina siliquis exuta denotant.*

5 . *Terebinth. minoris æmula, folijs angustioribus, & rotundioribus, Arbuscula Bisnagarica.*

6 . *Teucrij facie Bisnagarica tetracoccos rostrata, pilis scatens, folijs profundè venosis.*



Tab. LXV.

F. 1. *Teucrium lucidum parvo folio flore venustè purpureo. Chamædrys Alpina lucida* Cat. H. R. P. *Chamædrys Alpina latifolia elatior splendens* Sch. Bot. *Teucrium African. fruticans folio viridi splendido* D. Magnol. Mus. Courten.

2. *Teucrium Atticum molli lanugine incanum, an Teucrium Creticum* Clus. Hist. Park. *Teucrium Creticum incanum* C.B.P. *fortè etiam Scordium lanuginosum ejusd* C.B.P. *Scordium lanuginos Creticum* Camer. *munificentiâ* D. Wheeler. *accepimus*.

3. *Thalictrum pratense majus Monspeliensium foliis rugosis* Cat. H. R. P.

4. *Thalictrum minimum fœtidiss.* Caspare Bauhino I B. Hist. Park.

5. *Thalictrum minus lucidum Libanotidis Coronariæ foliis, an Thalictr. foliis lucid. Rorismarini instàr quorundam.*

6. *Thalictr. angustiss. folio* Park. *Thalictr. pratense angustiss. folio* C.B.P. *quod prim. in Prodr.*

7. *Thapsia latifolia Hispanica* Park. *Thaps.* 4. *Clusii. latifolia flore luteo, semine lato similis Sesedi Peloponensi* Lobelio I B. Hist. l. 3.

8. *Tithymalus s. Chamæsyce altera virginiana foliis crenatis & maculâ fuscâ eleganter notatis, noster,* Raii Hist. Commelin. Hort. Amst. *triginta abhinc adminimum annis in Horto* D. Walker. *observavimus.*



Tab. LXVI.

f. 1. *Thapsia glutinosa segmentis densioribus subhirsutis, alterno situ positis. an Seseli Peloponense folio Cicutæ Adv. & Icon. Lob.*

2. *Thapsia Cæsalpini semine candido craſſo I B. Hist. t. 3.*

3. *Thaps. latifol. villosa C. B. P. Seseli Peloponens. majus Lob. Ic. Thapsia quorund. hirsuta & aspera, Cicutæ folio, flore luteo, semine lato, alijs Seseli Peloponeſiacum I B. Hist.*

4. *Thapsia glutinosa Pimpinellæ Saxifragæ folijs venosis. ex Hispaniâ Thapsia 4.ª s. Tuero. dicta Clus. Hist. Thapsia 4.ª Clusij. Tuero. vocata folijs glutinosis, radice nigrâ foris I B. Hist. t. 3. Thapsia folijs Libanotidis fœtidiſsima C. B. P. item Cicutaria latifolia fœtidiſsima altera Phytopin.*

5. *Thapsia Carotæ facie I B. Hist. Thapsia Carotæ folio C. B. P. Thaps. vulgaris Carotæ effigie Adv. & Ic. Lob. lit. a. hujus umbellam seminibus onustam indicat.*



Tab. LXVII.

f. 1. *Thapsia maximis folijs glutinosa, segmentis densioribus ex adverso sitis. Thapsia Salmanticensis. s. 4ª. Clusij. magna, flore luteo semine lato* I B. *Hist. t. 3. Thapsia maxima Hispanica* Park. *Thapsia maxima latiſsimo folio* C. B. P.

2. *Thapsia Libanotidis folio glutinosa glabra. Thapsia Thalictri folio* Bot. Monsp. *Thapsia s. Turbith Garganicum, semine latiſsimo* I B. *Hist. t. 3. lit. a, appingitur Umbella.*

3. *Thymelæa Africana glabra floribus rubérrimis nostra* Ray. Hist. app. *an Sanamunda 4ª. s. folijs glabris acutis* Breyn. Fascic.


4. *Tribulus terrestris Americanus Argemones flore flavo, Tribulus terrestris major Curaſsavic;* P. B. P. *Tribul; terrestris Americ, Chamæcisti flore, fructu spinoso maximo* Breyn. Prod. 2. *lit. a, fructum denotat, ab Insigni Botanico D. Tancred. Robinson, accepimus, qui in Hortis Batavis collegit, sed absq; nomine.*

5. *Trifolium purpureum Ebenus Cretica dictum, . Loto affinis folio & facie Pentaphylloides fruticosi nostratis, floribus in spicam longiorem dispositis. Cytisus Creticus* ἀϱχονδόξυλον. Honor. Belli.

TAB. LXVIII.



Tab. LXVIII.
f. 1. Trifolium siliquos Loto affine, siliquis Ornithopodij. Trifolium siliquis Ornithopodij nostras Raÿ Synops. & Hist.
2/2. Trifolium pediculis foliorum longiſsimis Bisnagaricum, fortè Trifolij Bituminosi genus Indiæ Orientalis.
3. Tulipifera Caroliniana folijs productioribus magis angulosis Mus. Courten. fortè Macpalxochis-Quahuitl Mexicensibus Hernandez. Hist. lit. a. florem expansum indigitat.
4. Tulipifera virginiana Laurinis folijs aversâ parte rore cæruleo tinctis, Conibaccifera. Laurus Tulipifera baccis rubellis Cono erecto inclusis. Banist. Cat. Laurus Tulipifera folijs subtus ex cinereo aut argenteo purpurantibus Ray Hist. lit. b. baccas Cono capsulato per tenue filamentum appensas ostendit.



Tab LXIX.
f. 1. *Trifolium pumilum Bisnagaritum argenteo nitore sericeum capitulis oblongis hispidis.*
2. *Trifolium Americanum elatius holosericeum summo folio bifido suaveolens. Meliloti genus (fortè) Indianum.*
3. *Vitis Idææ species Maderaspatana Niruri fortè Malabarensibus dicta* Hort. Mal. P. 2. *an Frutex Indicus baccifer Vitis Idææ secundæ Clusij facie* Breyn. Cent. 1.
4. *Vitis Idæa Bengalensis Uvæ Ursi baccâ singulari, ad foliorum exortum barbulis donata.*
5. *Volubilis Gangetica sapore acri, & aromaticâ, clarius autem Aristolochia Clematitis Gangetica folijs in aculeum abeuntibus, radice odoratissimâ, an Aristolochia Clematitis Indica odorata* P. B. P. Addend. *Carelee-Vengo* Hort. Malab. *Sasanda Zeylanensib.*
6. *Volubilis, s. Aristolochia Clematitis, Bisnagarica, spinis gemellis aculeata.*
7. *Wadoukæ Malabaricæ haud multum dispar, Frutex aculeatus è Maderaspatan* Hort. Mal. P. 4.

TAB. LXX.



Tab. LXX.

f. 1. Verbena Scutellariæ, s. Caſsidæ folio, diſpermos, Americana, àn Verbena Indica Bontij. Hist. Ind. Or. fortè etiàm Verbena Curaſſavica Scutellariæ folijs, flore purpuraſcente P. B. P.

2. Veronica Virginiana procerior, folijs ternis, quaternis, & etiàm quinis, caulem amplexantibus, spicis florum candidiſsimis. àn Veronica Americana erecta, frutescens, & ramosa folio Verbenæ P. B. P.

TAB. LXXI.




Fig. 1. Fig. 2.

Tab LXXI.

f. 1. *Vicia multiflora maxima perennis tetro odore floribus albentibus, lineis cæruleis, Striatis, Vicia Sylvatica multiflora maxima* Phytol. Brit. Ray. Synops & Hist. *Vicia perennis multiflora spicata major, floribus albis, lineis cæruleis notatis*. Moris. Hist. 2.

2. *Vicia Lathyroides nostras, s. Lathyrus Viciæ formis* Ray Hist. & Synops. *Lathyrus ex cæruleo & rubro mixtus* Merret. Pin.



Tab. LXXII.
f. 1. *Vicia multiflora major angustifolia glabra, floribus oblongis ex albo & cæruleo mixtis. an Vicia Onobrychidis flore C.B.P. fortè Vicia maritima multiflora alba Meſsanensis* Ray Hist. & Cat. Exter. *fortè etiam Aracus minor Lusitanicus* Park. *Legumen pallidū Ulyſsipponense* Nonij. Brãdonij. Ger. emac. App.
2. *Vicia multiflora Caſsubica frutescens Lentis siliquâ* Breyn. Prodr. 2. Cat. Hort. Ac. Lugd.

LEONARDI PLUKENETIJ

PHYTOGRAPHIA

Seu

Stirpium Εἰκωνες fideliter & exquisite ad vivum delineatæ ÆNEISq TABULIS illustratæ

in Quibus

Novæ plures & Exoticæ Plantæ à nemine hactenus animadversæ nunc primum proponuntur. Necnon aliæ quædam apud Neotericos dubiæ atq obscuræ (mutatis aliquandò Nominibus prout res ipsa postulat) perspicuâ brevitate dilucidantur.

PARS ALTERA

ET SIC MEMINISSE IUVABIT



I Collins fec:

Ea quæ scimus, sunt pars minima eorum, quæ Ignoram.

LONDINI. MDCXCI.

SUMPTIBUS AUTORIS.

INCLYTO HEROI

atq;

ILLUSTRISSIMO DOMINO.

DOMINO

à BENTINGK.

COMITI de PORTLANDIA. &c. &c &c.

φιλοβοτανῶν PATRONO longè Gratioſiſſimo.

LEONARDUS PLUKENETIUS

Hortorum amænitates, ac adumbratam VEGETANTIS ORBIS. Imaginem, Hisce TABULIS suis PHYTOGRAPHICIS in altera Parte contentis, exhibitam, summâ cum humilitate.

Dat. Dicat. Dedicat.



Tab. LXXIII.

f. 1. *Absinthium Ponticum repens vel supinum C.B.P. Absinth: Ponticum supinum Herbariorum Obs. Lob. Absinth. repens Tab. Icon.*

2. *Absinth. Halepens. tenuifolium grati odoris, comâ delicatiori. Absinth. Santonicum Arabicum D. Bobart. fortè Scheha Arabum, Absinthij genus circa Bethlehem Rawolfy. Hist. Lugd. App. Hac pulcherrima Absinthij specie, cum plurimis alijs novis & Illustrioribus Plantis quas summâ diligentiâ ac indefeſſis peregrinationibus, per totam ferè Europam, sibi conquisivit Ornatiſſimus vir, & hujus Botanices Scientiæ verè Ornamentum D. Gulielm. Sherard. nos ditavit.*

3. *Acacia spinosa foliolis perexiguis, & veluti crispatis Maderaspatana, fortè Frutex folijs Acaciæ Mesopotamic. Fab. Col. Not. in Recch. Acaciæ similis Mesopotamica minutiſſimis folijs, siliquâ integrâ contortâ, craſsâ, & obtusâ, s. Siliquâ Nabathæa Breyn. Prodr. 2.*

4. *Acaciæ affinis virginiana spinosa, siliquâ membranaceâ planâ, florib. albis, papilionac. Anagyrid. modo in Uvam propendentibus. Acacia virginiana siliquis glabris Ray Hist. Arbor siliquos. virginiens. spinosa, Locus Nostratibus dicta Park. Anagyris Americ. spinosa floribus albis odoratis P.B.P. item Laburnum odorat. Cartadens folio Coluteæ. ejusd. P.B.P. Laburnum spinosum Mauritanicum folijs Glycyrrizæ flore albo Breyn. Prodr. 1. Glycyrrhiza, vel si mavis, Glycyrrhizæ affinis spinosa, arborescens Mauritanica, Laburni flore albo, ejusd. Br. Prodr. 2. lit. a. siliquam indigitat.*

5. *Adianthum exoticum Rutæmurariæ æmulum, pinnulis oblongis, angustis, in summitate bipartitis, à D. Pardeo habuimus.*

6. *Adianth. maritimum segmentis rotundioribus, Sch. Bot. Adianth. mas Tab. Ic. C.B.P. ubi planè fatetur, quid sit hoc Adianthi genus, se nondum aſſequi. Jerseiæ Angliæ nostræ adjacentis Insulæ alumna est, ut candidè nos monuit. D. Sherard.*

7. *Altea an Malva potiùs exotica, Pentaphylli folia profundè secta, purpureis floribus sine pediculis.*



Tab. LXXIV.
f. 1. Alcea Indica frutescens in lacinias variè diſsectis folijs. an Alcea arboreſcens Japonica folio Quercus, flore pleno carneo, in medio roseo; Cl. Breyn. Prodr. 2. Sansanqua Japonensibus.
2. Alsine Saxifraga Graminifolia flosculis tetrapetalis herbidis & muscosis, an Saxifraga Anglica annua Alsine folio D. Plot. Ray Hist. fortè etiam Saxifraga graminea pusilla folijs brevioribus craſsiorib; & succulentioribus ejusd. Synops. Septentrionalibus Angliæ nostræ a D. Lawson observata.
3. Alsine Polygonoides folijs tenuiſsimis per genicula binis ad summum caulium polyanthos. Alsine Nodosa multiflora Sch. Bot. a nunquam satis laudato viro D. Sherard. habuimus.
4. Alsine palustris repens folijs lanceolatis floribus albis perexiguis. Alsine palustris exigua folijs lanceolatis, Plantaginellæ aquaticæ instar, flosculis albis vix conspicuis Mentz. pugil. Plantago aquat. minim. Clus. Hist. Park. Plantaginella palustris C. B. P.
5. Alsine aquatica major repens, folijs acuminatis, Virginiana.
6. Althæa Indica Abutili subrotundo folio, flore luteo, spicato, fortè Malva Americ. Abutili folio, flore luteo spicato, folijs incanis P. B. P.
7. Althæa Scammonij folio, flosculis albis Zeylanica P. B. P. Nilavonanghu & Wiſadulu Zeylanensib; dicta in Hort. Regio. Hamptoniens. longe celeberrimo, conspeximus, & in Horto Comptoniano collegimus.
8. Althæa Americana fruteſcens, Melochiæ folijs angustiorib; an Malva frutesc. Melochiæ angusto folio, florib; albis fasciculatim congestis P. B. P. inter Addend. an Tsjeru-Uren. Hort. Malab. P. 9.



Tab. LXXV.

f. 1. Alsine montana, capillaceo folio. C.B.P. Park. an Alpina Saſsifraga angustifolia, minima montana Col. ecphr. quæ parum differt ab Alsine tenui-foliâ muscosâ ejusd C.B.P.

2. Alsine Caryophylloides angustifolia, multiflora, glabra, purpurascens, radice Astragaliti, an Caryophyllus sylvestris, repens, multiflorus C.B.P. forte etiam Caryophyllus Saxatilis foliis gramineis, umbellatis corymbis, ejusd C.B.P. (i.e) Symphytum petræum, s. Gypsophyton majus Thaly. Harcy. nid

3. Alsine Polygonoides tenuifolia, flosculis ad longitudinem caulis veluti in spicam dispositis nostra. Ray Synops. Polygonum angustiſsimo gramineo folio, erectum. Bot. Monsp. in Maritimis circa Bostoniam agri Lincolniës.

4. Alsine Polygonoides herbacea minor, Lariceis foliis capillaceis, ex uno pediculo plurimis. à D. Sherard. habui.

5. Alsine Alpina glabra foliolis reflexis. an Saxifraga Bavarica Park. ejusd D. Sherard. humanitate accepi.

6. Alsine Spergulæ facie Bengalensis, foliis angustis ad genicula binis, flosculis autem plurimis, ad caulem radiatis, semine minutiſsimo fulvo.


7. Alsine Americana Nummulariæ foliis Parad. Batav. Prodr. ab eodem D. Sherard. communicata.

8. Amaranthus Indicus verticillatus albus, Origani foliis lanugine incanis. an Amaranth. Alopecuroides Indicus herbaceus, Atriplicis folio. Breyn. Prod. 2 forte etiam Amaranthoides humile Curaſsavicum foliis Polygoni Par. Bat. Pr. in Hort. Reg. Hampton. observavimus

9. Amaranthoides humile Curaſsavicum foliis Cepææ lucidis, capitulis albis. P.B.P. Amarantho affinis aizoides, s. Amaranthoides minor Americana procumbens, Sedi teretifolii foliis, & facie, flore oblongo niveo. Breyn. Prod. 2. Trifolii spicâ Crithmum maritimum non spinosum Braſiliense Ray. Hist. Perexil. do Mar. Lusitanis. s. Caaponga 1ª Braſiliensium Pison. Hist.



Tab LXXVI.
f. 1. Angelica Charibbæarum elatior, Olusatri folia flore albo, seminib; striatis, Cumini odore & sapore.
2. Anchusa repens Lithospermi facie, floribus cæruleis, secundum folia prorumpentibus. Lithospermum virgatum Thaly. Harcyn. Lithosp. maj; repens latifolium C.B.P. Lithosp. majus Dodonæi fl. purpur. semine Anchusæ I B. Hist. t.3. l.33. p.592. ubi Icon minus feliciter exsculpta, per errorem ad pag. 692. transposita est.
3. Anchusæ degeneris facie Indiæ Orientalis, herba quadricapsularis.
4. Anonis purpurea procumbens verna, s. præcox, fruticosa, annua, viscosa, siliquis craſsis, lentiformibus, varius dispositis. Moris. Hist. 2. fortè Anonis Hispanica frutescens, folio Ciceris sylvestris triphylli, Sch. Bot. Anonis annua ramosa, folijs rotundis serratis Cat. Hort. Edinburg.
5. Apocynum hirsutum scandens Americanum flore amplo purpureo, an Apocynum scandens Africanum hirsutum P.B.P. & Spectatiſsimi viri D. Fosteri Horto plurimarum Plantarum exoticarum instructiſsimo, Lambethæ ad Austrum sita florentem vidimus & habuimus.
6. Apocynum Curaſsavicum fibrosâ radice florib; aurantijs, Chamænerij folijs angustioribus P.B.P. an Apocyn. erect. minimum angustif. Americ. flore umbellato, aurantio, petalis reflexis Breyn. Prod. 2. sed nostri florum petala non reflexa sunt, sed potius inflectuntur, & ità disposita, ut flosculum concavum efforment.



Tab LXXVII.

f. 1. Apocynum American. erectum tuberosâ radice non incanum, folijs rigidioribus, latis, subrotundis, florib; alb. intùs purpureis summo cauli Corymbum magnum efformantibus. an Apocyn. Americ. arborescens, folijs rigidis, subrotundis, ampliſsimis. P. B. P. fortè Apocynum erectum non ramoſum, latiore folio, Umbellis florum albicantibus Banister. Cat. Stirp. Virg.

2. Arbor Americana Morifolia, vasculo seminali verrucoso, Coni æmulo, ovali, Balsamum olente. Cedrus spuria Jamaicensib; nostratibus dicta.

3. Arbor Orientalis Bellidis spinosæ solidioribus folijs, capsulis parvis quadripartitis, Kiluke Cingalensib; dicta hujus Specimen exsiccatum Insigni Botanico D. Tancredo Robinson. debemus.

4. Arbor Americana triphylla, fructu Platani quodammodò æmulante. Lignum fibularium (i.e) Buttonwood, Nostratib; dicta.

5. Arum triphyllum minus pene atro rubente Virginianum. Arisarum triphyllum minus pene atro rubente Banister.

6. Arum Moschatum Ophioglossoid. s. pene nudo Jamaicense. an meliùs Ophioglosſum Moschatum Jamaicense Limonij folijs, in extremo Sinuatis. àn Ariſarum fluitans pene nudo Virginianum. Banist. Cat. MSS.

TAB. LXXVIII.



Tab. LXXVIII.
f. 1. *Aristolochia polyrrhizos, auriculatis folijs, Virginiana, fortè Polyrrhizos Virginiana Park. Serpentaria altera Virginiana vulgò.*
2. *Asarum Virginianum Pistolochiæ folijs subrotundis Cyclaminum more maculatis. Asarum Cyclamini folio Virginiana. Banist. Catal.*
3. *Asparagus aculeatus triplice spina Prom. Bon. Spei. an Asparagus sylvestr. spinosa Africana Cat. Hort. Amst. colitur in Hort. Reg. Hamptoniensi*
4. *Aster luteus Cisti hirsutis, & crenatis folijs, summo cauleramoso, floribus amplis. an Herba Vulneraria. ad fluvium Pfrim. Hieronym Tragi. Conyzæ affinis Germanica C. B. P. Consolida Saracenica 3.a Tragi. Hist. Lugd. Conyzis affinis, Herba Vulneraria, s. Solidago Saracenica 3.a Trago hirsuta I. B. Hist. t. 2. hujus exemplar nobis amicissime donavit D. Doody. qui & ipse inter Siccatas exoticas, & quamplurimas rariores Plantas, in Horto Potavino jamdudùm collectas, & in Chartis asservatas, quas sibi impetravit, absq; titulo invenit.*
5. *Aster Americanus Belvidere folius, floribus ex cœruleo. albicantibus. spicis longioribus. an Aster Novæ Angliæ Tripolij floribus spicis longis dispositis P. B. P.*
6. *Aster American; multiflorus, flore albo Bellidis, disco luteo D.nœ de Flines. Sch. Bot.*



Tab. LXXIX.
f. 1. Aſter American; albus latifolius, caule ad ſummum brachiato, ſi color florum, & foliorum conjugata poſitio non obſtarent, hanc Tlaloteloxochitl Nov. Hiſp. Recch. eandem plantam eſſe ſuſpicaremur.
2. Aſter Americanus albus Mexerei Arabum ex aſperatis folijs, florum petalis reflexis.
3. Aſtragalus annuus folijs & ſiliquis hirſutis, plurimis in foliorum alis ſeſſilibus. Aſtrag. ſiliq. & folijs hirſutis floribus parvis Moris. Hiſt. 2. Fœnu=græco ſylveſtri Tragi in quibuſdam accedens Planta I B. Hiſt. t. 2. Aſtragalus marinus Bœticus Park. Vicia Seſamacea Apula Col. ecphr.
4. Aſtragalus hirſutus corniculis plurimis uno pediculo ſtelli formiter adaptis. Ornithopodio affinis hirſuta fructu ſtellato C.B. P. Stella leguminoſa I B. Hiſt. t. 2. munere Ornatiſſimi Viri D. Patricij Adair accepimus.
5. Aſtragalus Pyrenaicus incanus, folijs Viciæ, flore ſulphureo. Sch. Bot.
6. Auricula Urſi Virginiana floribus albis Borraginis inſtar roſtratis, Cyclaminum more reflexis. à Celeberrimo Botanico D. Baniſter. in Angliam nuperrime tranſmiſſa.



Tab. LXXX.
f. 1. Balsamum Caprarium Antegoanum v. Nevisense fœtidum. s. Arbuscula Americana Balloteolens, Scammoneæ foliis punctatis, ferrugineâ lanugine obductis, floribus parvis pentapetalis spicatis.
2. Braſsica spuria exilis non laciniata, caule magis folioso hirsutior, foliis subrotundis. Braſsica spuria minima, caule magis folioso hirsutior nostra. Ray. Synops. app.
3. Brunella Hyſsopifolia C. B. P. Brunella minor folio rigido, integro, ejusd. Phytop. Prunella angustifol. I B.
4. Bugula non crenata tomentosa, Suecica, seminibus albis. munere doctiſs. viri & Divini Verbi Prædicatoris apud Londinates Reverendiſs. D. G. Stonestreet. prorsus novum hoc Bugulæ genus accepimus.
5. Buxus Africana, rotundifolia serrata P. B. P. in Horto Regio Hamptoniens. collegimus.
6. Buxus Lauri Alexandrinæ foliis accedens Americana.
7. Cardiaca minor annua, Americana, flore argenteo. s. Leonurus Americana alba, folio sideritidis longiore. è seminibus natam, superiori anno, in Horto Comptoniano observavimus.



Tab. LXXXI.
f. 1. *Caſsia lignea Jamaicensis, cortice acri candicante. Canella alba Clusij. Laurifolia Magellanica cortice acri C.B.P. Cortex Winteranus acris, s. Canella alba I B. an Ligni aromatici Monardis cortex.*
2. *Carduus Tingitanus, flore magno, cæruleo, folio Attractylidis divisurâ subtùs incano, spinis durioribus horrido.*
3. *Caryophyllus sylvestris Alsines holosteæ arvensis glabræ folijs flore unico, calyce barbato Armeriæ species flore summo caule singulari nostra Raij. Synops. app. forte Caryophyllus sylvestr. humilis flore unico C.B.P. an Lychnis Meſsanensis parvo flore rubello, e calyce oblongo angusto Raij. cat. Extorav.*
4. *Centaurium ciliare minus, Bisnagarium, Origani folijs amplioribus, floribus in Umbellis.*
5. *Chamædryfolia, Scandens claviculata & Corymbifera Jamaicensis. huc fortè pertinet Bryonoides. s. Temecatl-Yauctepecensis Mexicensibus Hernand.*




Tab. LXXXII.
f. 1. Bellis montana minor Tanaceti folijs Moris. Prælud. Tanacetum montan. inodorū, minori flore C. B. P. Tanacetum album I. B. Hist. I. 3. Tanacetum inodorum secund. Clus. Hist.
2. Centaurium Ciliare minus, Circææ folijs firmioribus, spicatis floribus è Maderaspatàn.
3. Ceratia diphyllos Antegoana, Ricini majoris fructu ossea siliquâ grandi incluso. Arbor siliquosa ex Virginia lobo fusco scabro. C. B. P. itèm Arbor siliquosa ex quâ Gummi Anime elicitur. ejusd. C. B. P. Jetaiba arbor Gummi Anime fundens. Pison. Acaciæ quodammodò accedens, Arbor Anime Gummi fundens, Americana folijs magnis, acuminatis, in pediculo binis, Lobo magno crassissimo, eduli. Breyn. Prodr. 2. Animifera arbor Brasiliana P. B. P. hujus siliqua est Lobus ille ex Wigandecaoro, de quo Clusius. in exot. Locus Nostratib; bulgò. aa. Lobum. bb. bb. fructum indicant.
4. Ceratiæ quodamodo affinis, Bengalensis, folijs bigemellis, subrotundis, siliquis admodùm intortis, & in orbes circumflexis, ex minio nigricantibus fructu rubro, maculâ nigrâ insignito. àn Aoaramo-Temo Brasiliensibus Pisonis? a. siliquam b. fructum indigitant. in Horto Comptoniano enutritam vidimus.

TAB. LXXXIII.



Tab. LXXXIII.
f. 1. Chryſanthemum minus Alpinum elegans, flore magno albicante foliis craſsis, Saxifragiæ chelidonides accedentib; hæc elegans planta Chryſan-themo Corymbifero Africano Triumfetti, quoad folia, optime convenit, a D. Parry, ad exsculpendam accepimus.
2. Chryſanth. Orellanæ foliis impense scabris Maderaspatanum, florum petalis quinis tetragonis, ex calyce amplo viridi fistuloſis.
3. Chryſanth. anguloſis Platani foliis Virginianum. Doronicum maxim. American. latiſsim. anguloſis fol. radice transparenti Cat. H. Ac. Lugd. an Chryſanth. latifol. Braſilianum C.B. prodr. forte Noviytepuzt quadrilatera Hernandez.
4. Chryſanth. Virginian. villoſum disco luteo petalis quinis ornato. Chryſanthem. pentapetalon villoſo caule Banist. Cat. Mſs.
5. Chryſanth. hirſutum Virginian. auriculato Dulcamaræ folio octopetalon. Chryſanth. folio oblongo auriculato Banist. Cat. Mſs.
6. Ciſtus humilis. Chamæciſtus Ericæ folio luteus elatior C. B. P. Helianthemum tenuifol. glabrum flore luteo I B. Hist. t. 2. Chamæciſtus Ericæ folio Park. Cori ſimillimus, Frutex in maritimis Andr. Cæſalp.



Tab. LXXXIV.
1. Chryſanth. Alpin. Bellidis folio caule singulari. Chryſanth. parvum, s. Bellis lutea parva latifolia I B. Hist. t. 3. Chrys. Alpestre Bellidis folio Cat. H. Pat. à Turre.
2. Chryſanth. Alpinum Bellidis folio ramosum.
3. Chryſanth. Americ. bidens alatum flore parvo aurantiaco. Chrys. conyzoides alato caule Curaſſavicum P.B. P. Chrys. cannabinum American. alatum, flore aphyllo, globoso, aurantio, folijs Baccharidis Breyn. Prodrom. 2.
4. Cistus Helianthemos Ericæ brevioribus folijs & humilior. C. B. P.
5. Cistus Alpina humilis, folijs Thymi minutiſſimis.
6. Cist. humilis Maſſiolotica Camphoratæ tenuiſſimis folijs glabris Chamæcist. Maſſiliens. Corruda vel Coridis luteæ folio Schol. Bot. hanc eandem eſſe plantam, cujus ôia folia tenuiſſima Camphoratæ similia & glabra, quam Celeberrim; noster Rajus circa Maſſiliam invenit. ab ornatiſſimo Botanico D. Sherard. meritò censetur.
7. Clematis Alpina Geranifolia C. B. P. & Prodr. Clematis cruciata Alpina Pona. Descripl. Montis Baldi.
8. Clinopodium minus exoticum Thymi folio majore. Clinopod. Creticum Pr. Alpin. Thymum exoticum flosculis nunquam expansis D. Bobart.
9. Conferva geniculata minima nostras. Oculatiſſimus D. Sherard. hanc eſſe Umbelliferam aliquando suspicabatur.



Tab. LXXXV.
f. 1. *Clinopodium minus exoticum Thymi folio minore. Clinopodium Thymi folio & facie, Schol. Botan. Has 5que plantas noviſsime recensitas, à D. Sherard accepimus.*
2. *Clinopodium Amaraci folio, flore albo Americanum Dictamnus ibi vocatum.*
3. *Convolvulus Sagittariæ folijs, flore purpureo Wheeleri. ab Inventore habui.*
4. *Convolvulus Scammonea dictus macrorhizos. an Scammonea macrorhizos Pr. Alp. exot. Scammonea macrorhizos. Cretica Park. hanc enutrivit jampridem in Horto suo D. Edvard. Morgan.*
5. *Cucumis parva repens Virginian. fructu minimo Banist. Cat. MSs. c fructum d. d. semina denotant.*
6. *Cupreſsus Virginiana folijs Acaciæ corniger æ paribus & deciduis, Cupreſsus Virginiana folijs Abietis mollib; atq deciduis Breyn. Pr. 2. Cupreſs. Americana Park. Tlatzan. f. Cupreſsus Americana fusa Hernand. an Habhel. Paludanus cujus fructus seu formam species seu squammarum amplitudinem ab hujus nuce haud multum abludit, ut recte notat Cl. Rajus; ob loci tamen Natalis distantiam eandem non ausus est affirmare. Ita. a. a. hujus nucem integram & diversimode diſsectam. b b. nucleos ostendunt.*



Tab LXXXVI.
f. 1. Cytisus foliis viridib; angustis quasi complicatis Monspeliens. C.B.P.
2. Cytisus procumbens Americanus, fl. luteo ramoſiſſimus; qui Anil suppeditat apud Barbadensium colonos. Chamæ-Indigo sylv. trifoliata Barbadensib; an Cytisus humilis procumbens P.B.P.
3. Doronicis affinis longifolia Ray Hist. Tuſsilago Alpina folio longo I.B. Hist. t. 3. Tuſsilago Alpina folio oblongo C.B.P. Doronicum Pyrænaic. Persicæ foliis glabrum Schol. Botan. ubi pro non descript. ponitur.
4. Elatines folio piloſiſſima papilionaceo flore Maderaspatana.
5. Eliochryso affinis Orientalis foliolis ad genicula binis, flore pentapetaloide argenteo apicibus aureis paleaceo.
6. Eliochryso affinis Oriental. ramosior foliolis caulium nodos radiatim ambientib; Comâ paleaceâ, flore albo lituris aureis striato.
7. Eruca sylvestris minor lutea Bursæ pastoris folio C.B.P. Eruca pumila Bursæ pastoris folio Boccon. Eruca Monensis laciniata lutea Raÿ Hist.
8. Eruca Virginiana Bellidis majoris folio Moris. Hist.



Tab. LXXXVII.
f. 1. Eupatoria conyzoides Verbaſci folia, s. Baccharis Americana ſuaveolens, Euedexylon (ie) Sweetwood. Barbadenſibus dicta.
2. Eupatoria conyzoid. odorata folio crenato, molli ſubincano. Serratula Noveboracenſis, folio leviter crenato, molli ſubincano P. B. P. fortè, Serratula Indica major latifolia mollis Breyn. Prod. 2. Iaceæ v Serratulæ affinis, capitulis Baccharidis, folijs Trachelij Zeylanica. ejusd. Prod. 1. Cattu-Schiragam Hort. Malab. P. 2. fortè, etiam Conyza Syriaca Rawolfij. Hist. Lugd. app.
3. Eupatorium Americanum, Meliſsæ folijs magis acuminatis. an Eupatorium Novæ Angliæ, Stachidis s. Betonicæ folijs crenatis, florib; albis P. B. P. fortè etiam Eupatorium Americanum Nepetæ folio latiore, floribus albis brevioribus; Breyn. Prod. 2.
4. Eupatoria conyzoides maxima, Canadenſis folijs caulem amplexantibus. Conyza major, alato caule, folijs Primulæ veris longè amplioribus, graviter odoratis. Lælij Triumfetti. apud Fratrem. an Virga aurea latiſsimo folio, Canadenſis glabra Schol. Botan.
5. Eupatoria Conyzoid. integro Iacobææ folio glabro v duriuscule, Indiæ Orientalis. fortè Iaceæ v Serratulæ affinis Breynij. Cat. H. Ac. Lugd. app. hanc enim Cl. Hermann: à Cattu-Schiragam. Hort. Malab. non ſeparat, cum verè verè ſunt Plantæ inter ſe admodum diverſæ, ut ex Iconibus noſtris liquido patebit. Eruditiſsimo Dn. Sherardo hujus notitia debetur.
6. Eupatorium Virginian. Salviæ folijs longiſs. acuminatis perfoliatum. Huic folia longa mucronata, bina oppoſita, v ad baſin ità continua invicem, ut ea caulis transadigatur, ſinum quendam, ſeu cavitatem pelvi aquæ continendæ ſimilem, efformant.



Tab. LXXXVIII.

f. 1. Eupatorium humile Africanum, Senecionis facie, folijs Lamij P.B.P. Senecio Africanus Lamij folio Acad. Reg. Paris. Senecio Americanus Lamij facie Schol. Botan. fortè Virga aurea humilis Alpina Moris. Præludd. fortè etiam, Virga aurea folio Veronicæ Mentzelij. Faſcic.


2. Eupatoria hirſuta Hyſſopifoliorum æmula Virginiana, an Tlalchinol-Xochitl s. Venerea Medicina 2ª. Hernandez Hiſt. Mexic. p. 238. & 415.

3. Eupatoria Valerianoides, flore niveo, Teucrij folijs, cum pediculis, Americana.

4. Eupatoria Valerianoides, Triſsaginis folio, absq; pediculis, Virginiana, Cacalia folijs rotundioribus, ad caulem ſeſsilibus Banister. Cat. Mſs. an Tlacoxochitl Nov. Hispaniæ Recch. p. 418.

5. Evonymus Canellæ albæ folijs è Maderaspatan, Dris. Frederici Slare favore, ad exsculpendam accepimus.

6. Evonymo adfinis Arbor Orientalis nucifera flore roseo. Styraci & Evonymo media affinis, The Sinensium, sive Tsja Iaponensibus flore simplici, Breyn. Prod. 2. Thee frutex Bontij hujus flores munere D. Petiver accepi.



Tab. LXXXIX.
f. 1. Ebenus Jamaicensis Arbuscula Buxi folijs spinosa flore luteo papilionaceo siliquifera.
2. Filix saxatilis, non ramosa, minima. Ray. Cat. Exter. ar. ubi dubium movet an sit Dryopteris Dalechampij IB.
3. Filix saxatilis, omnium minima elegantiſſima. Cat. Hort. Reg. Paris.
4. Filix Alpina Myrrhidis facie Cambrobritanica an Filicula fontana Adianti facie latifolia dentata Schol. Botan.
5. Filicula pumila Lonchitidis Maranthæ species Cambrobritannica, an Lonchitis aspera Ilvensis Hist. Lugd.
6. Filix parva Cambrobritannica pinnulis latis, auriculatis, brevibus, aculeata. has tres noviſſimè enumeratas Filices ab Inventore D. Lloyd. accepit insignis Botanicus D. Sherard. a cujus liberalitate & ipse habui.
7. Filix minor Africana Lonchitidis folia, pinnulis auriculatis planis. Filix Africana minor Lonchitidis facie P. B. P. in Hort. Regio. Hamptoniens. longè celeberrimo ᵉ collegimus.
8. Filix Virginiana pinnulis ablongis, angustis, auriculatis, tenuiſſimè crenatis, longè ab invicem distantibus, non spinosa.
9. Filici-folia s. Polypodium tenuifolium minus Virginianum.
10. Fagi folijs alterno positu alatis, Arbuscula Bisnagarica.



Tab. XC.
f. 1. *Fœnum græcum Phaseoloides, Virginianum, flore amplo cæruleo. Securidaca trifolia flore patulo cæruleo, radice Glycyrrhizæ sapore Banister Cat. Mſs.*
2. *Fumaria Tingitana, radice fibroſa, perennis, flore ex albo flaveſcente, ſiliquis curtis. an Fumaria semper virens & florens ex Hortis Batavis Cat. H.R.P.*
3. *Fumaria ſiliquoſa, radice grumoſa, flore bicorporeo, ad labia conjuncto, Virginiana. Fumaria siliquosa, grumosâ radice flore gemello monstroso. Banister Cat. Mſs. an Fumaria Canadensis, radice tuberosâ squammatâ Acad. Reg. Paris.*
4. *Fungis congener, Acetabulum petræum, s. Fungus minimus petræus, orbiculatâ crustulâ compositus. D. Lloyd. Ray Synops.*
5. *Galegæ degener, folio singulari, floribus parvis herbidis muscosis. Galega unifolia, flore herbaceo. Cat. Hort. Palav. Georg. à Turre.*
6. *Gentiana Maderaspatensis, foliis Æsclepiadis, caule ramoso.*



Tab XCI.

f. 1. Geniſtella Montis ventoſi ſpinoſa I.B. Hiſt. t. 1. Geniſtella Monſpeliaca ſpinoſa C.B. prodr.

2. Gramen cyperoides exiguum, ſeminibus bullatis, Raij Obſervat. 444. ex mente D. Sherard. qui nobis ad exſculpend. dedit.

3. Gram. Cyperoid. Americanum, ſpicis ſpecioſis plurimis, compreſſis, ſummo caule abſq; pediculis adnatis. an Gramen maritimum Cyperoid. Curaſſavicum panicula punicante P.B.P. Gram. Cyperoid. minus, panicula ſparsa ſpadicea. fortè Jacape Braſilienſibus Piſonis.

4. Gram. Cyperoid American. ſpicis majorib; oblongo rotundis Sparganij in modum echinatis, ad ſummū caulem pediculis longis innitentibus.

5. Gram. dactylon Orientale, majus frumentaceum, ſemine Napi. Coracan Cingalenſibus D. Knox. Itinerār. p. xi.

6. Gram. parvum Gangeticum, ſpicâ gemellâ habitiori. an Gram. δυσαχυόφορον. Col. ecphr. 338. hujus ſpicæ Amaraci capitula ſquammata quodammodo exprimunt.

7. Gram. Cyperoid. folijs Caryophylleis, ſpicâ granis rarioribus, & tumidioribus conflatâ Raij Synops.

8. Gr. Cyperoid. tenuifolium, ſpicis ad ſummum caulem ſeſſilibus, globulorum æmulis. Gr. Cyp. folijs mollib; tenuibus ſpicis brevibus & coacervatis, tribus quatuorve in ſummitate caulis. Raij Hiſt. fortè Gr. Cyp. minimum, Ranunculi capitulo rotundo D. Bobart. ejuſd. Raij Synops. Stirp. Britan.



Tab. XCII.

1. Gram. dactylon Siculum multiplici paniculâ, spicis ab eodem exortu geminis Ray Hiſt. an Feſtuca junceo folio ſpicâ geminâ C. B. P.
2. Gram. amoris alterum, paniculis ſtrigoſioribus magiſq; ſparſis Ray, Cat. Exter. gr. in Hiſtoria autem omiſſa est.
3. Gram. American. spicâ echinatâ majoribus locuſtis Schol. Bolan. fortè etiam Gram. aculeatum Curaſſavicum P. B. P. ſeminibus à D. Sherard. communicatis in Horto D. Doody, ex oticarum refertiſſimo enatum primò obſervavimus.
4. Gram. Alopecuroides Indicum minus ſpicâ Sericeâ candidâ laxiori.
5. Gr. Indicum Alopecuroid. holoſericeum majus, ſpicâ longâ pappeſcente. an Gram. Alopecuroides Curaſſavic. ſpicâ longiſſimâ Sericeâ candidâ. P. B. P.
6. Gramen Miliaceum Americanum minus paniculâ parvâ.
7. Gr. Miliaceum American. maximum, paniculâ minore.
8. Gr. Miliaceum Americanum medium, paniculâ magis ſparsâ & ſpecioſâ.
9. Gram. junceum elatius, pericarpijs ovatis Americanum.



Tab. XCIII.

1. Hedera Clematitis Atragenes singulari folio splendente Virginiana.
2. Helleborine Virginiana, Ophioglossi folio. D. Banister. hujus Exemplar pulchrè exsiccatum, nec non picturam propriâ manu ad vivum apprimè delineatam Eruditissimus Inventor nonitâ pridem amicis suis in Angliam transmisit.
3. Hieracium folijs in tenues lacinias, profundè sectis, flore luteo. Chondrylla lutea I B. t. 2. Chondrilla Coronopi, aut Cyani facie, tenuifolia lutea binedrum Adv. Sonchus lævis in plurimas lacinias divisus C.B. Pin. & Prodr.
4. Hypericum ferè orbiculato folio floribus amplis, pallidè luteis, petalis florum in ambitu crenatis. Hypericum Nummulariæ folio C. B. prod.
5. Hypericum Ericoides minimum, folijs cinereis, ex Hispaniâ, an Hypericum Francorum, non scriptum, tenuifolium Belluc. Ind. Hort. Pisan. Invenit in Hispaniâ D.r Turnefort.
6. Hypericum Alpinum, magno flore, nigris punctis insignito Sch. Bot. fortè Ascyron magno flore C. B. Prodr. Has tres proximè antecedentes, novas, aut saltèm, minùs cognitas Hypericorum species, munere præstantissimi viri D. Sherard accepimus.
7. Jacea tenuifolia purpurea, capitulo Calcitrapæ in modum aculeato. Capitulis spinosis Jacea Adv. Jacea cum spinosis capitulis, purpurea, tenuifolia I B. t. 3. lib. 25. an, Stoebe squammis asperis C. B. P.

TAB. XCIV.



Tab. XCIV.
f. 1 Jacea marina Boetica Park. an Jacea purpur. maritim. capitulo spinoso Neopolitana Raij Cat. Exter.
2. Jacea peregrina Napifolia hirsutis capitulis alata. e seminibus a D. Sherard, cum ab Exteris in Angliam rediret, secum allatis, et ab ipso communicatis, in Hortis quibusdam Curiosorum, apud Nostrates nata est, et nuper viriles succrevit in Hort. Acad. Oxon. sub auspicijs Dni Bobart. qui hujus specimen nitidè exsiccatum nobis amicissimè transmisit.
3 Jacea fruticans Pini folio C. B. P. Stoebe capitata Rosmarina folio Ponæ, s. Chamæpitys fruticosa Cretica Belli. I B. t. 3. Stoebe fruticosa angustifolia Cretica Park. an Chamæpeuce Pr. Alpin. exot. 77. Illustrioris hujus Plantæ nobile specimen a Reverendiss. D. Wheeler. ad æsculpend. impetravimus.
4. Jasminum vulgò American. at rectius Evonymo adfinis Occidentalis, alatis Rusci folijs nucifera, Cortice ad genicula fungoso. Guaiacum American. alterum fructu Evonymi Breyn. Prod. 1. Guaiacum folijs Lentisci Horto Fageliano. ejusd Breyn. Prodr. 2. Packhout Belgarum sub Jasmini titulo. in Hort. Reg. Hamptoniens. primùm nobis innotuit. a. fructum occlusum, quatuor obtusis angulis constantem, ostendit.



Tab. XCV.

f. 1. Laureola latifolia floribus albicantibus, odoratis Americana. sub nomine Jasmini Americ. Aurantiæ folio collegimus in Hort. Reg. Hamptoniens.

2. Imperatoria Ranunculoides Africana, enneaphyllos. Laſerpitij lobatis folijs rigidis, margine ſpinoſis. Ranunculus per elegans D. Franqueville, folijs aculeatis Lob. Illustr. dt. 135. Sphondylij ꝛ. Panacis rigido, hirtoq; folio, Planta Africana caustica. P. B. P. Nostra tamen folia sunt penitùs glabra, & lævore splendentia. ideò q; tota planta Laſerpitio propius accedit, quam Sphondylio, Umbellifera urens. Cap. Bon. Spei. bulgo. in Hort. Reg. Hampton. Obſervavimus.

3. Juncus Orientalis, paniculâ ſpecioſâ è Maderaspatan. Humaniſſimi Viri & Mercatoris Londinensis celeberrimi D. Hileard. favore accepimus.

4. Kali Curaſſavicum procumbens, folijs subrotundis P. B. P. 344.

5. Kaka-Toddali (fortè) Malabarica; ex Oris Cormandel. H. M. part. 5. Frutex baccifer. Indicus, ſpinoſus, trifolius, floribus ſpicatis, fructu pleno, rotundo, tricocco Raij Hist.

6. Limonium (fortè) peregrinum, folio ſolidiori tecturâ in lacinias pariè diſſecto. à D. Edv. Morgan. jam olim exſiccatum accepi.

7. Juncus elegantissimus capitulis pectinatis Prom. Bon. Spei. D. Sherard amicè communicabit



Tab XCVI. 1. Laserpitium exoticum, lobis longiſsimis integris Ammeos quorundam diviſuris, à D. Sherard habui.

2. Liguſticum humilius Scoticum à Maritimis: s. Apium maritimum dulce Scoticum. Apium Scoticum & Apium marinum quibusd. Newlono. Raij Hiſt. Imperatoriæ affinis Umbellifera maritima Scotica Sibbald. Prodr. & Cat. Hort. Edinburg. an Liguſticum quorund. foliis Angelicæ I B. I. 3. quod in Alpib. Liguriæ bulgò Imperatoria bocatur, ut in hujus Deſcript. notat Cæſalpinus. an Libanotis Apii folio ſemine aſpero C. B. P. Roſmarinus foliis Selino ſimilibus, quæ Libanotis legitima Theophr. Belli. epiſt. 1. ad Cluſ. σκοτιοχεα bulgò, quod bertiginem inducere credatur, & ni fallor (inquit Bellus) Siler Cretic. à quibusd. perperàm bocatur. Siler marit. Scoticum Imperatoriæ facie Breyn. Prodr. 2.


3. Linaria lutea flore minimo C. B. P.

4. Linar. triphylla major floribus parvis pallescentibus, rictu aureo, calcari subcæruleo, donatis. an Linaria latifolia, triphylla major. C. B. P. Linaria Cretica latifolia Cluſ. Hiſt.

5. Loto affinis Coryli folio Acad. Reg. Paris. Anthyllis Ulmifolia Quorundam.

6. Lycium Italicum Bod. à Stap. in Theophr. Lycium Buxifolio C. B. P. Lycium Buxifoliis rotundioribus, Syriacum, bel Perſicum Breyn. Prod. 2. Osrum Perſarum, Rhamni species Adolph. Vorſt. Cat. Hort. Acad. Lugd.

7. Laurifolia fructu parvo, typhoide non papposo, gemello, ſiliquas æmulante, Arbuscula Maderaspatana.



Tab. XCVII.

1. Lycium minus cordatis Buxi folijs Maderaspatanum. forte Lycium Hispanic. folijs Buxi C. B. P. Tamaios. Lycium verum Fragos. an Lycium Buxi folijs angustioribus Syriacum. Breyn. Prodr. 2. Lycium Dioscoridis. Hadhad Arabibus, & Zaroa Incolis Montis Libani. Rawolfij. Itinerar.

2. Lycium Maderaspatanum, Indici Alpino putati æmulum bijugis, & grandioribus aculeis horridum.

3. Lycium Bisnagaricum, acuminatis, minus durioribus folijs, & aculeis ex opposito binis. Tsjerou-Karæ Hort. Malab. simile, si non idem.

4. Lycium putatum Indiæ Orient. Capparis rotundioribus folijs, & aculeis ex adverso gemellis.

5. Lycium Beloxylon, binis aculeis in ramulorum fastigijs, Americanum. ex hujus ligno sagittas, e fructubus Atramentum conficiunt Indigenæ. Dartwood & Inke-berry à Nostratibus nuncupatur, à D. Sherard. specimen ad exsculpendum datum est.

6. Lycium majus Americanum, Iasmini flore, folijs subrotundis lucidis. Iasminum spinosum Americanum folio lucido subrotundo P. B. P.

7. Lycium Myrti folijs subrotundis, Americanum, lactescens, limbis foliorum argentatis.

8. Lycium (fortè) non spinosum, crassis & incanis folijs P. B. P. rectius autem, Cerasus Sebestenæ domesticæ folijs, aliquatenus accedens. Prom. Bon. Spei. Capsi Carsiboom vulgò Belgarum.


9. Lychnidis, aut potius Alsines cognata dispermos; s. Chamæjasme tetrapetaloides inodora, quadrato caule Virginiana.

TAB. XCVIII.



Tab. XCVIII.
f. 1. Lychnidea umbellifera Blattariæ accedens Virginiana major, repens Pseudo-melanthij folijs pilosis, flore albo, pentapetaloide, fistuloso.
2. Lychnidea Blattariæ accedens Virginiana minor, repens, hirsutis Camphoratæ folijs, floribus fistulosis, in Umbellam positis.
3. Lychnidea Blattariæ accedens Virginiana Camphoratæ congeneris glabris folijs, flore ad summitatem ramulorum singulari.
4. Lysimachiæ purpureæ affinis Americana procumbens, Anonidis bernæ frutescentis folio singulari glabro, Gratiolæ affinis frutescens Americana folijs Ageratoidis s. Veronicæ erectæ majoris Breyn. Prodr. 2. hanc plantam Capræ apud Belgas Curassavienses pro pabulo avidè petunt inde Capraria Curassavica, Cabritta vulgo P. B. P. quam D. Wheeler sponte natam in Grecia Observavit, ubi Cusermobt à Turcis nuncupatur. Florem habet nivei coloris pentapetaloidem, capsulam cylindricâ figurâ bipartitò divisam seminibus purvis turgidam.
5. Lysimachiæ purp. affin. fruticosior, Anonidis singulari hirsuto folio Maderaspatensis.
6. Malus Granata Zeylanens aculeata. Granata Malus Zeylanica spinosa P. B. P. an Kalu-naregam H. M. part. 4. in Horto Reg. Hampton. collegimus.
7. Melilotus Indiæ Orient. minima dispermos folliculis in summitate aduncis an Melilotus minima, Syriaca Nephel, s. Nephal Ibnbaithar, & Malasesæ Breyn. Prodr. 2.
8. Muscus polytrichoides pellucidus fere, folijs denticulatis, ad margines buli crispis noster. Ray Synops. Stirp. Britan. append. an Polytrichum aureum medium ejusd. Ray Histor.
9. Muscus denticulatus minor C. B. P. Park. Muscus terrestris Lu. ... I B t. 7.

TAB. XCIX.



Tab. XCIX.

f. 1. *Marrubium aquatic. fol. integris longis per angustis Maderaspatanum. àn Marrubium palustre Canadense tenuiùs glabrum Sch. Bot.*

2. *Melampyrum cristatum I. B. t. 3. àn Melampyr. luteum angustifol. C. B. P. Park. quærit & ipse Cl. Rajus.*

3. *Melampyro affinis tetraphylla Gangetica, floribus inter folia sparsis, àn Adel-Odagàm H. M. part. 9. hujus semen, mellei coloris lucidum, singulare, nudum, grani tritici æmulum insidet super basi foliorum cristatorum duplici serie binas membranulas rigidiores, alteram alteri longiorem, circum ambientium constructa.*

4. *Mercurialis tricoccos hermophroditica. s. ad foliorum juncturas, ex foliolis cristatis julifera simul & fructum ferens Banist. Cat. Stirp. Virg. à Mercurial. Zeyl. tricocco, cum acetabulis Cat. H. Ac. Lugd. append. plurimùm differt, ut ex utriusq. picturâ, cuivis pateat.*

5. *Mespilus Virginiana spinis longioribus rectis, folijs quodammodò auriculatis.*

6. *Millefol. palustre, galericulatum minus. Reverend. Vir & amicus noster singularis D. Dodsworth. in agro Eboracensi sponte natum, invenit & jampridèm nobis benignè communicavit.*



Tab. C.
f. 1. *Meſpilus Virginiana Groſſulariæ foliis.*
2. *Muſcus rupeſtris, repens, Virginianus, clavis folioſis, erectis, quadratis Baniſter. Cat. Mſs.*
3. *Muſcus filicinus elegans, Bonæ ſpei Prom. Bon. Sp. Putturjoory Hottentottorum à D. Sherard. accepimus.*
4. *Myrrhis annua ſemine roſtrato, Striato, aſpero oblongo nodoſa Moriſon. Umb. Myrrhis ſylveſtr. Cretic. nodoſa ſeminibus aſperis Ray. Hiſt. Seſeli Cretenſe nodoſum Park. Chærophyllon alterum, geniculis tumentibus. Cat. H. R. P.*
5. *Myrrhis perennis lutea Daucoides Moris. Umb. & Prælud.*
6. *Myrtus Brabanticæ affinis Americana foliorum laciniis Aſplenii modo diviſis, julifera ſimul & fructum ferens. fortè, Canambaya Braſilienſib. Marcgrav. fortè etiam, Cardiaca Mulleriana Indica Bod. à Stap. in Theophr. 320. hujus ſemina ſingularia capſulis bivalvib. extrà globum echinatum emergentibus, continentur.*
7. *Eadem, cum ramulo conis, s. nucamentis, onuſto.*



Tab. CI.
f. 1. Muscus palustris in Ericetis nascens floridus, noster. Ray Synops. App. : at minus incautè floridum pronunciavim; capitula quidem hujus, pedicellis innixa, versus ramulorum apices, operculis eorum jam delapsis, flosculos tam pulchrè æmulantur, ut primo intuitu nobis imposuere; ex quo demum agnosco, nihil aliud eſse quam Muscus palustris terrestris similis Ray Histor. Pl.

2. Nardus Americana procerior folijs cæsijs, huic proxime cognata est Teripeti bel Tzipalitzili Nov. Hisp. Ant. Recch. p. 412.

3. Nasturtium petræum Cambrobritannicum Iohnson Merc. Bot. part. alt. Paſsim in excelsis rupibus Arvoniæ & Mervinjæ v. g. Moelyn rhudh juxta Phestiniog, y Clogwin, du y yr Ardhu, Clogwin y Garnedh prope Lhandberys invenit Plantarum Britannicarum ſagaciſsimus Investigator D. Lloyd.

4. Nastort. Bursæ pastoris folio Virginianum flore albo, siliquâ compreſsâ in Hort. bene Urbano D. Doody primo Observavimus.

5. Nastort. African. floribus albis spicatis, & folijs ternis Christophorianæ facie P. B. P. in Hort. Reg. Hamptonicas. collegimus.

6. Nil. s. Anil sylvestre Cynanchicæ folijs Bisnagaricum. an Manneli Hort. Malab. P. 9. quam Genistellæ tinctoriæ speciem facit Commelinus in Not. supr. loc.

7. Ninzin, s. Zingin, & Ginseng Radix genuina Indiæ Orientalis, apud Miſcell. Curios. Ephem. Germanorum. 2. Ann. 51. p. 73. sub hujus titulo plurimas Icones videsis, inter quas Ginseng Cleyeri Iaponiensis Terræ alumna huic nostræ quam maxime accedit. Quum Ninzin Bontij ab hac planta plurimum differt. frondium effigiem ope & favore Cl. Viri Dris Tancredi Robinson. hujus radicem munere Amiciſsimi Viri D. Gedley accepimus;



Tab. CII.
f. 1. Onobrychis Maderaspatana diphyllos siliculis asperis. hujus notitia, humaniſs. aequè ac Clariſs. Viri D^vis^ Frederici Slare favori debemus.
2. Onobrychis, s. Hedysarum Clypeatum Alpin. perenne siliqua glabra, fl. albo Sch. Bot. Onobrych. Clypeata laevis C. B. P. & in Prod. à D. Sherard. habuimus.
3. Ornithogalo affinis Virginian. flore purpureo pentapetaloide Banist. Cat. Mſs.
4. Oxys purpurea Virginiana, radice Lilij more nucleata, capitulis postquàm defloruerit bulbillis, ut in Allio corvino conflatis. Trifolium acetoſiſsimum, radice parva rotunda Lilij more imbricata Banist. Cat. Mſs. a. capitulum bulbillis suis onustum in digitat.
5. Pedicularis. s. Cristae galli affinis Virginiana Ajuga multifido folio, apicibus coccineo floribus pallidis in spicam congestis. Euphrasia Virginiana comis coccineis flore albido theca foliaceâ incluso Banister. Cat. Mſs.
6. Phaseol; tuberos; Cap. Bon. Spei sub Batatas Capensis trifoliatae titulo accepimus.
7. Potomogeito similis Graminifolia ramosa, & ad genicula polyceratos. Potomogeito affinis Graminifolia aquatica Ray. Histor. Plantar.



Tab. CIII.
f. 1. Palma Hayri Park. Theatr. Palmæ facie Hajri I B. l. 2. Palma American. spinosa C. B. P. Palma Braſiliens. vi. s. Ayri Piſon. p. 129. Palmiſte epineus Rochfortij p. 60. forte Macow Arbor Barbadenſium Ligonis. hujus Arboris lignum marmori nigro simile, ferri modo in aquâ fundum petit, exinde Ebeni speciem eſse quidam autumant. plurimas hasce ex Insulâ Jamaicensi alldtas in Horto Comptoniano aliquando conspeximus.

2. Palma American. folijs Polygonati brevioribus, leviter serratis, & nonnihil spinosis, trunco craſso Breyn. Prod. 2. Palma Americana non spinosa, craſsis rigidisq; folijs caudice scabro in caput protuberante P. B. P. in Horto. Reg. Hampton. bidimus & collegimus.

3. Periclymeno recto accedens Arbuscula Virginiana siliquosa.

4. Pimpinella Sanguiſorba rupestris minima, spinosa radice magna. hujus folia Pimpinellæ, foliolis haud majora, à Cl. Dre. Turnefortio in Hispan. collecta fuit, & à D. Sherard amiciſsimè nobis communicata est.

5. Plantago Coronopus dicta, maritima, brevi, latoq; & dentato folio pilosa. Coronopus Maſsiliensis hirsutior latifolius Sch. Bot. Coronopus folijs acutis in margine dentatis Ray Hist. Pl. Plantagini affinis Bibinella, Sicilis herbula I B. l. 3. Coronopus maritimus Prochytæ Insulæ Col. ecphr. 258. non autem est Coronopus maritima nostras, nec ei multum similis ut Rerum omnium Naturalium, Vir peritiſsimus, & Botanices Scientiæ Summus Arbiter & Præsidium singulare, D. Ioh. Rajus rectè pronunciat. Cat. Exterar. quamvis Casp. Bauhinus eandem faciat. Plantago Apula bulbosa Park. Plantago Apula laciniata bulbosa Fab. Column. Plantago pilosa bulbosa C. B. P. huic nostræ persimilis est & fortè eadem.

6. Polygonum Germanicum caule, folioso, flosculis in summitate. an Polygonum anguſtiſs. & acuto vel gramineo folio montanum Sch. Bot. fortè Polygono anguſtiſsimo folio affinis C. B. P. s. Hæmorrhoidalis quæ ad Caryophylleorum sylvestrium claſsem referri debere Clusius putavit. Hist. app. alterâ. ad. lib. 6. fortè etiam Polygonum Polonicum cocciferum I B. l. 3.



'Tab. CIV.

f. 1. Paſsiflora Vesicaria hederacea, folijs lanuginosis, odore tetro, filamentis florum ex albo & purpureo variegatis. Hedera Murucuice species Vesicaria lanuginosa, odoris gravis Marcgr. 73. Flos Paſsionis Curaſsavicus, Hederæ hirsuto folio, flore purpureo variegato P.B.P.

2. Periclymenum herbaceum rectum Virginianum. D.[r] *TinKarsweed. ibi vulgò vocatum hujus generis eſse videtur illa Periclymeni species scandens, & corticibus Arborum radices emittens, Itti-Canni* Hort. Malab. *dicta.* part. 7.

3. Phaseolo affinis Arbor Indica Coral dicta polyphyllos, non spinosa, folijs mollibus subhirsutis.

4. Piperi longo similis Pistolochiæ folijs absq. pediculis Maderaspatana. Jaborandi 4.[a] *Pisonis huic multùm convenire videtur* Hist Brasil. p. 216.


TAB. CV.




Fig. 1. Fig. 2. Fig. 3. Fig. 4. Fig. 5. Fig. 6.

Tab. Cv.

1. *Phaseolus unifolius Occidentalis Asari benoso folio dispermos Orobus Americanus Asari folio singulari* P. B. P. *in Hort. Regio. Hampton. enutritum observavimus.*
2. *Phaseolus unifolius Orientalis summo folio sinuato monospermos. àn Phaseolus minimus perennis Americanus flore luteo Meliloti; folijs Asari lanuginosis solitarijs Breyn. Prodr. 2.*
3. *Phyllitis parva Saxatilis, Virginiana per summitates foliorum radicosa Banist. Cat. Stirp. Virg. Herba derata. hujus folium Phyllitidis ramosæ Alpini folium particulare figurâ æmulatur et huic forsàn ista proximè cognata est, alterius, ramificationis defectum alteri per admi=rabilem Propogandi modum sibi ferè peculiarem suppeditante beneficâ Naturâ.*
4. *Portulaca Curassavica lanuginosa Kali folio erectior, & elatior flore rubente* P. B. P. *Portulaca Americ. erecta lanuginosa folijs Psyllij, florib; in caulium summo rubro obsoletis majorib; Breyn. Prodr. 2.*
5. *Portulaca lanuginosa procumbens Vermiculatæ folijs Americana. Portulaca Americana procumbens Sedi teretifolij folijs, & facie floribus in ramorum summo pulchrè purpurascente Br. Prodr. 2. Beldroga fortè Lusitanis, s. Caaponga Brasiliana 1.a Pisonis 244.*
6. *Portulacæ affinis Americana, tricapsularis erecta flore albo, pediculis angulosis. Portulac. Americ. s. Curassavic. latifol. erect. florib; albis* P. B. P. *ex hujus capsulâ tripartitò divisâ semina per maturitatem impetu dissiliunt. has tres novissime enumeratas plantas in Horto. Regio. Hampton. oculis nostris summâ voluptate lustravimus atq; collegimus.*



Tab. CVI.
f. 1. *Phœnico-Scorpiuros, s. Heliotropium Palmites spinosum. Polygonati angustifolijs Maderaspat. huc forte pertinet Atitara Braſilienſibus Marcgr. lib. 2. p. 64.*
2. *Ejusd. summitas floribus ornata, quæ nodosas & incurvatos Scorpionum caudas pulchrè repræſentat, ut in Heliotropio. favore Cla. D. Frederici Slare habuimus.*
3. *Ranunculus Nemorum Fragariæ folijs Virginianus. Banister: Cat. Mſs.*
4. *Ranunculus Nemorosus Aquilegiæ folijs, Virginian. Asphodeli radice. Ranunculus Thalictri folijs grumosâ radice Banister. Cat. Stirp. Virg. nuper edit.*
5. *Ranunculo adfinis Umbelliferis accedens, Chelidonij minoris folio Zeylanica minor. Asarina minor Zeylanica P. B. P.*
6. *Ranunculo adfinis Umbelliferis accedens Chelidonij minoris folio, Curaſſavica major Asarina major Curaſſavica P. B. P.*
7. *Pentaphyllum Alpinum minus folijs tenuioribus altius serratis, cauliculo purpuraſcente. an Pentaphyll. repens minus aureo flore Clus. Hist. cui folia magis villosa attribuit, nostra equidem vix hirsuta apparent. hujus exemplar amiciſſimus Vir & apud Londinenses Pharmacopæus peritiſſimus D. Philippus Schrickenfauchs nobis conceſſit.*



Tab. CVII.

f. 1. *Pyrolæ affinis Virginiana, repens, frutiosa, folijs rigidis Scabritie asperatis, flore pentapetaloide fistuloso. Pyrola repens folijs scabris, flore pentapetaloide fistuloso.* Banister.

2. *Reseda Linariæ folio Pyrenaica glauco folio* Sch. Bot. *an Reseda Linariæ folio* C. B. P. *item Reseda alba minor ejusd.* C. B. P.

3. *Rhamnus Maderaspatanus Evonymi pallentib; folijs, spinis brevibus duplicatis.*

4. *Rhoi similis Tragôodes, leptiphyllos, Americana, spinosa, rachi medio appendicibus aucto. an Coriaria Arbor spinosa Acaciæ folijs & facie* P. B. P. *inter* Addend. *forte Quauhtecopalli Pitzahuac, Copallifera leptiphylla alia* Hernand. p. 49. *Fingrego, & Sabina arbor (i.e.) Savin Tree Barbadensib; nostratibus dicta.*


5. *Ricinus parvus urens, folijs Quercinis Virginianus* Banist. Cat. Stirp. Virg. *an Quercinis folijs Arbuscula spinosa lactescens Porto ricenſis* P. B. P.

6. *Ricini fructu parvo, frutiosa, Curaſsavica, folio Phylli,* P. B. P.




Tab. CVIII.
f. 1. Rhamnus Maderaspatanus Hypericifrutescentis folio, spinis brevibus duplicatis.
2. Rhamnus an potius Lycium Fingrego Jamaicensibus dictum, forte Katu-Karu-Walli H. M. spinarum autem positio huic parum congruit.
3. Ricinus Maderaspatanus quinquifidis durioribus foliis in margine spinosis, humanitate & favore D.ris Fred. Slare Obtinuimus.
4. Rubus foliis eleganter diſsectis D. Fagon. in Hort. Reg. Paris. collegit D. Sherard & nobis communicavit.
5. Sanicula Alpin. anguſtiſs. folio, flore carneo. Sedum Alpin. alterum Franciſc. Gregory. Regiensis Cappucini. Fab. Column. ecphr. p. 2. Sedum Alpinum anguſtiſsimo folio, flore carneo C. B. P.
6. Sanicula Alpina anguſtiſsimis brevioribus foliolis per intervalla confertis. Sedum Alpin. primum Column. ecphr. p. 2. Sedum Alp. exiguis foliis C. B. P. an Sedum Alpinum minimum carneo flore, Sch. Bot.
7. Scabiosæ capitulo, Chrysanthemi Myconi foliis, alato caule Maderaspatensis.

TAB. CIX.



Tab. CIX.
f. 1. Scabiosa Conyzoid. Americana capitulis, & floribus albidis parvis P. B. P.
2. Schetti album, s. Iasminum Indicum Laurifol. inodorum, umbellatum floribus albicantibus P. B. P Bem-Schetti. Hort. Malab. P. 2.
3. Serratula Noveboracensis maxima foliis Doriæ leviter serratis P. B. P. inter Addend. Serratula Virginian. min. angustif. mollis Breyn. Prod. 2. forte Atzoyatl Cozticyni-Roderilyo. Nov. Hisp. Recch. 355. huic congener est Tlalli-yetl Nov. Hisp. ejusd. pag. 423.
4. Sesamum alterum foliis trifidis Orientale, semine obscuro. in Car-Elu Hort. Mal. P. 9. Bidajam Kitern s. Bidajam nigrum Malaccensibus, Sesamum Indicum, folio amplo serrato, flore majore, semine nigricante Commelin. Annot. in Hort. Malab.



Tab. CX.

f. 1. Scutellaria Teucrij facie purpurea minor, flore ampliſsimo. Teucrium Astragaloides Caſsida minor Cat. H. R. P. 173.

2. Scutellaria major, Nepetæ folijs Virginiana. an Scutellaria aquatic. major Americ. Cat. H. Ac. Lugd. fortè Tochoneochitl. Nov. Hisp.=
=Ant. Recch. 431.

3. Smilax Viticulis aspera folijs longis; angustis; mucronatis lævib; auriculis ad basim rotundioribus.

4. Smilax Viticulis asperis Virginiana, folijs angustis, lævibus nullis auriculis prædita huic radix geniculata craſsa & intorta, indè suspicamur Chi=
=næ Mexicanæ speciem eſse, & fortè Olcacatzan Hernand. p. 212. & Ioapecanga altera Pison. fortè etiam Radix Chinæ Acost. Hist. Lugd. app. 10.

5. Smilax Virginiana spinis innocuis armata, latis Canellæ folijs radice arundinaceâ craſsâ & carnosâ. fortè China spuria nodosa C. B. P. Pseudo-
-China radix ex Virginia Clusij. exot. China Michoacanensibus Phaco dicta Hernand. 213. Pocayo Nov. Hisp. Ant. Recch. 398. Ipape=
=canga Brasiliensib; Pison. 258. an Kari-Vilandi H. M. P. 7. Pockhout Belgis Orientalib; utibidem habetur. apud nostrates Virginiam
habitantes, ex hujus Stipitib; parantur Scipiones, ut nobis affirmârit Ampliſsimus & Senatoriâ Dignitate perhonorandus Vir D. Byrd. Utraq.
viz hæc simul & proximè præcedens Ioepecangarum species, fructum acinosum ac racematim dependentem in summis caulibus copiosè proferat.

6. Smilax aspera Bermudensis grandiorib; folijs cordiformibus, radice surculosâ. fortè Quautimecapatli. s. Zarzæ species cujus radices in Hispan=
=iam transferri solebant ut à Francisco Ximene narrat Marcgravius p. 12. Quauhmecapatli altera, Zarza 4a. Hernand.
290. forte etiam Smilax aspera Zacynthi Alpino I. B. t. 2.



Tab. CXI.

f. 1. *Smilax folijs latis in margine ſpinoſis Caroliniana, Stipite leni quadrato. hanc Generoſiſſimus Dominus Courtene nobis benigniſſime conceſſit.*

2. *Smilax viticulis asperis Virginiana folio hederaceo leni, Zarza nobiliſſima nobis. Mecapatli s. Zarza-parilla Fr. Hernand. 288. in Hort. instructiſſimo D. Foſteri invenimus.*

3. *Smilax Caroliniana, Stipite quadrato leni folijs angustis, asperis, auriculis ad basim angulosis. ab illustri viro D. Gulielm. Courtene accepimus.*

4. *Solanum Americanum Strychnodendro accedens fructu medio (ante maturitatem) quaſi ſſo. an Solan. frutesc. Braſilian. folio Capsici, baccis rubris P. B. P.*

5. *Solan. Curaſſavic. vesicarium, Solano Antiquorum simile folijs Origani subincanis P. B. P. forte Solan. vesicarium Surinamense lanuginosum polycarpon Br. Prod. 2.*

6. *Solanum triphyllum flore tripetalo atro-purpureo, in foliorum sinu, absq. pediculo Seſsili. Banist. Catal. Stirp. Virgin. Hujus autem florem nobis videtur, sex foliolis sive petalis constare, quorum tria subjecta tota viridia sunt, ubrevia, tria vero superiora, longiora, atro-purpurea sunt, quemad-modum in Solano triphyllo Canadensi Cornuti. Cui hæc est quam proxime cognata.*

TAB. CXII.



Tab. CXII.
f. 1. Senecio American. altiſsim. Blattariæ vel Hieracij folio Sch. Botan. Senecio Americ. maxim. Dipsaci s. Virgæ Paſtoris folio P. B. P. an Senecio Cichorei folio Virginiana flore sulphureo odore Paſtinacæ Cat. Mſs. H. Oxon.
2. Solan. Barbadense erectmosum minus tinctorium Circææ folijs mollib. & incanis forte Heliotropium Curaſsavicum Scammonij folijs mollib. subhirsutis. P. B. P. Heliotropium Americanum majus folijs Populi nigræ lanuginosis flore albo minimo. Breyn. Prodr. 2.
3. Spongia fluviatilis anfractuosa perfragilis ramoſiſsima nostras. in Isi fluvio prope Oxonium invenit D. Sherard & nobis munere donavit.
4. Spongia marina fistulosa maxima muricata. an Spongia ramosa fistulosa. C. B. P. dedit nobis D. Doody.
5. Syringa Volubilis Virginiana Myrti majoris folio, alato semine, florib. odoratis, luteis. Gelseminum s. Jasminum luteum odoratum Ray Hist. Gelsemin. s. Jesmin. Virginian. scandens & semper virens Park. Theatr. Pseudo-Apocynum Myrti-folium Virginian. Banist. Cat. Stirp. Virg.
6. Tithymalus Curaſsavic. Salicis & Atriplicis folijs varijs, caulib. viridantib. P. B. P. Tithymalus Indicus, Salicis & Atriplicis folijs Breyn. Prodr. 2.



Tab. CXIII.
f. 1. Tithymal. exotic. Carobe foliis aliquatenus accedens. an Tithymalus Siculus polycoccos Portulacæ folio Bocc. 39. Tithym. baccifer Hyacinth Ambros. Phytolog. 532.
2. Tithym Indic. annuus dulcis, floribus albis, cauliculis viridantibus P. B. P.
3. Thymelæa Africana Linariæ folio, florib. albis in extremo caulium odoratiſsimis. Thymel. Lini foliæ similis African. florib. pallidis odoratiſsimis P. B. P.
4. Trifolium stellatum glabrum Ger. emac. Trifol. capitulis Dipsaci quorundam Raÿ. Synops.
5. Trifol. supinum cum glomerulis ad caulium nodos orbiculatis florib. purpurantib. Raÿ. Histor. an Trifol. parvum rectum flore glomerato cum unguiculis I B. t. 2. meritò autèm quæratur (inquit eruditiſs. noster Rajus) cur rectum dicat, cum in Descriptione caulem supinum ei attribuat.
6. Trifolium cochleatum modiolis spinosis Raÿ. Synops. Stirp. Britan.
7. Valerianella Curaſsavica semine aspero viscoso P. B. P. in Hort. Reg. Hamptoniens. legimus.
8. Trifolium siliquosum Loto affine Maderaspatan. siliquis longis, ex adverso binis propendentibus.



Tab. CXIV.
1. Verbasco congener, è Maderaspatàn, mucronatis folijs, aversâ parte lanuginosis, flore parvo purpureo.
2. Veronica adfinis Indica tenuifolia, flosculis & caliculis barbatis, summo caulium fere umbellatis. a. vb. capsulam seminalem per Microscopium visam exhibet. è Maderaspatàn ad Ornatiſsimum virum D. Hileard allata est, qui nobis humaniſsimè communicavit.
3. Veronica saxatilis parva caulibus nudis Raij. Obs. pag. 442. munere D. Sherard accepimus.
4. Viburnum Americanum non spinosum Meliſsæ folijs floribus coccineis P. B. P. Camara flore rubro Marcgr. p. 5. fortè etiam Camara flore luteo ex ripa Fluvij S^{ti} Franciſci. ejusd. an Camara Juba Pison. Tbeiz quirh. s. Matzium torridum, & calidum Hernand. 115. nostri enim flores primùm lutei, deindè aurantij, tandèm miniati, & ultimò tanquàm coccinei sunt coloris, & in Hort. Reg. Hamptoniens. cum floribus & fructu conspeximus.
5. Viburn. Americ. odoratum folio parvo orbiculato, floribus & baccis folioli intercepta. P. B. P. 384.
6. Vicia Lathyroides s. Lathyrus Vicioides, vexillo rubro, labiab. petalis rostrum ambientibus cæruleis Moris. Hist. 2. Lathyrus s. Cicercula Meſsanensis Raij Hist.
7. Viola Virginiana tricolor, folijs multifidis, cauliculo aphyllo. Banist. Cat. Virgin. an Viola Alpina folio in plures partes diſsecta C. B. P. Viola montana laciniato folio Clus. Hist. Viola montana folio multifido I. B. Hist.
8. Virga aurea flore albo spicato. Breyn. Prod. 2. Virga aurea Noveboracenſis, floribus albis, folio Bellidis ramosa Canadenſis Cornuti P. B. P.

APPENDIX

ad Rariorum Plantarum Icones

LEONARDI PLUKENETY.

☞ Proptèr Literarij Sculptoris incuriam, aùt saltèm nostram in hoc opere properantiam contigit, Iconem Tab. 82. Fig. 5. adscriptam, absq; titulo exhiberi; hujus ergò Synonyma hìc subnectere lubet, ut suo loco reponantur. Viz.

5. Cerasa Hottentottorum; Lycium Africanum fructu Cerasi rubro. Parad. Batav. Prodr. 349. Rhamnus Africana Cerasi fructu ejusd. P. B. P. 369



Tab. CXV. f. 1. *Alni folia Americana serrata, florib. pentapetalis albis in spicam dispositis.*
2. *Ejusdem Spica florum, cum in majorem perfectionem devenerit.*
3. *Apocynum Virginianum Asclepiadis folio, erectum, floribus pallidè cæruleis, radice craſsâ. fortè Moyotlicyacatli Nov. Hispan. Ant. Recch. 388.*
4. *Chrysanthemum ex Insulis Charibeis Leucoij incanis, & sericeis folijs argenteis craſsis.*
5. *Evonymus Virginianus, Pyracanthæ folijs, semper virens, capsulâ verrucarium instar asperatâ rubente.*
6. *Fruticulus Americanus, cordatis folijs pentapetaloides, radice lignosâ.*
7. *Nymphææ congeneri Anapodophyllon dictæ accedens, polyphylla. s. Dracontij similis herba Americana pusilla.*



Tab. CXVI.
f. 1. Hieracium magnum Hiſpanicum Park. Dens Leonis latifolius caulescens C. B. P. & in Prodr. Hierac. Pyrenaicum, Blattariæ folio, minùs hirſutum, Sch. Bot. ubi pro nondum descripto ponitur.
2. Loto affinis hirſuta flore ſubrubente C. B. P. Anthyllis leguminoſa, purpuraſcente flore, Cluſ. Hist. Loto affinis hirſuta herba flore ſuave rubente. Moris. Hiſt. 2.
3. Methonica Malabarorum; Niengala Zeylanenſium Muſ. Zeyl. Cat. H. Ac. Lugd. app. Mendoni Hort. Malab. Lilium Zeylanicum, Superbum vulgò.
4. Thymus capitatus è Cithærá, foliis minimis confertis, an Thymus capitulis minoribus Maſsiolioticus Cat. H. R. P. à D. Wheeler habuimus.
5. Thymum majus longifolium, Stœchadis foliaceo capite purpuraſcente, piloſum, fortè Thymum capillaceo folio, hirſuto, magno capite purpuraſcente D. Turneforte. idem quod apud Collectaneas Rariorum quas Cl. Turnefortius ad D. Sherard è Luſitaniâ tranſmiſit, ſub hoc titulo aliquandò Conſpeximus.
6. Virga aurea Americana Tarraconis facie & ſapore, paniculâ ſpecioſiſſimâ, fortè Tlalchichinoa Patlahoal Mexicenſib. s. Herba uſta. 3.ª Hernand. an Oloxihuitl ejusd. Hernand. Hiſt. Mexic. de hujus sit genere determinare non audeo.
7. Fungus Virginianus ex Stercore Equino capillaceus, canus, capitulo rorido, nigro punctulo in ſummitate notato. Baniſter. Cat. Stirp. Virg. Mſs.



Tab. CXVII.

f. 1. Ficoides s. Mesembrianthemum Africanum frutescens, floribus albis parvis umbellatis. an Mesembrianthemum Afric. frutescens minus triangularibus foliis viridib. cornuum taurinorum in modum reflexis, fructu turbinato, parvo, pentagono, lignescente Breyn. Prodr. 2.

2. Rossolis Lusitanicus maximus foliis Asphodeli minoris D^ris. Turnefortii. Chamæleontoides Grisleij. in Epist. Dedicat. ad Viridar. Lusit. munere D. Courtene accepimus.

3. Serpentaria repens florib. Stamineis spicatis, Bryoniæ nigræ folio, ampliore, pingui, Virginiensis. huc etiam pertinet Mecaxochitl; Mexicensib. Hernand. 144. fortè etiam Xaxahoactli Nov. Hisp. Recch. 438.

4. Ejusd. spica flosculorum, cum ad justam explicationem assurgant.

5. Tulipifera Virginiana tripartito Aceris folio, mediâ laciniâ velut abscissâ Ray. Hist. Tulipifera arbor Virginiana Cat. H. Ac. Lugd. Populus, aut potius Alnus Tulipifera cornis erectis. Banist. Cat. Stirp. Virginiens.

6. Viscum arboreum flore albo specioso Americanum, foliis formâ siliquarum Nerij. Nerij siliquæ effigie, Kaye Bakal B. Hist. Fructus Nerij siliquæ figurâ C.B.P. per errorem.

ALTERA

APPENDIX.

ad Rariorum Plantarum Icones Quâ

Stirpes quasdam exoticas, proponere nos placuit, quarum Exemplaria ex India Orientali humaniſsimus vir, ac Chi= rurgus peritiſsimus D. Richard. Sambach *misit amico suo* D. Iacobo Pettever *Pharmacopæo, & apud Londinates Botanico meritiſsimo: à quo demum ad delineandas æreisq; typis exprimendas accepimus: Iſtis itaq; duobis cordatis viris grates agere de his debent qui studium Botanicum colunt*



Tab. CXVIII.
f. 1. Alſine, Spergulæ facie Maderaspatens. folijs latioribus semine minutiſsimo nigro. an Tzompili Huitzpatli Nov. Hisp. Recchij.
2. Amaranthus spicâ longâ candidiſsimâ, Spicæ folijs Maderaspatana.
3. Antirrhino affinis, pusilla planta, folijs scabris Maderaspatana.
4. Cardiaca s. Leonuros Hyſsopi hirsutis folijs Maderasp. floribus dilutè rubris, unico tantum verticillo in ramulorum cymis.
5. Chrysanthemum Maderaspatanum, Menthæ arvens. folio & facie, florib; bigemellis, ad foliorum alas, pediculis curtis.
6. Tithymalus botryoides, cordatis folijs, Indiæ Orientalis.



Tab. CXIX.

1. Gramen geniculatum, brevifolium, criſpum, ſpicâ purpuro-ſericeâ Maderaſpatanum.
2. Gram. dactylon Maderaſpatenſe, non pluribus ſpicarum paribus ad ſummum caulis, veluti in verticillas poſitis, folia magnâ ſui parte, culmum ceu vaginâ complectuntur, reliquam ſurrigunt breviorem, rigidam, marginibus criſpatam.
3. Gram. Cyperoid. Maderaſpatanum, Junceli Geſneri capitulis ſparſis.
4. Lyſimachiæ affinis Satureiæ folio Maderaſpatens. capſulis in ſummitate ferè umbellatis.
5. Melampyro cognata Maderaſpatana ſpinis horrida, an Coletta-Veetla, Hort. Malab. P. 9. Eryngium Zeylanicum febrifugum florib. luteis Herbar. Vivi. Herm. Cingalenſibus Kathu Karohiti, fortè etiam huc pertinet Bahel-Schulli Hort. Malab. p. 2. Geniſtæ ſpinoſæ ſpeciem facit, Commel. in Notis.
6. Phaſeolus arboreſcens incanus, monoſpermos, ſiliquis, lenti-formibus, Maderaſpatanus.
7. Sinapiſtrum Indicum diphyllon, ſiliquis ad alas ex uno pediculo binis. Cupa-Veela fortè Hort. Malab. P. 9.



Tab. CXX.
f. 1. Mimosa pumila Maderasp. pinnis tenuiſsimis, ad foliorum exortus innocuis spinulis obarmata.
2. Polygonum Solinoides Maderaspatens. Thymi capitati folijs incanum.
3. Portulacæ affinis Polygonoides, Bliti folio & facie Maderaspatensis.
4. Rhamnus cordatis folijs subtus cæſijs Maderaspatan.
5. Ricinus Althææ folio mollius incano Maderaspatens.
6. Ricinus urens Cannabinis folijs triphyllos è Maderaspatan. an Urtica Indica pentaphyllæa Caſsii Mich. Nutij. Catal.
7. Trifolium Maderaspatan. cauliculis pilosis scandens, Paſsifloræ modo trilobatum. an Coen. s. Coentic. Mexicensib; Hernand. 252. Apios Americana triphylla siliquis Orobi.
8. Viola surrecta Maderaspatensis Lini facie, rotundioribus imis folijs.